

© The Author(s) 2020. Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

STUDIES

Effect of aridity on species assembly in gypsum drylands: a response mediated by the soil affinity of species

Arantzazu L. Luzuriaga^{*1}, Pablo Ferrandis², Joel Flores³ and Adrián Escudero¹

¹Department of Biology and Geology, Rey Juan Carlos University, 28933, Móstoles, Madrid, Spain, ²Botanic Institute of UCLM, Botanic Garden of Castilla-La Mancha, Avda. de La Mancha, 02006 Albacete, Spain, ³División de Ciencias Ambientales, Instituto Potosino de Investigación Científica y Tecnológica, San Luis Potosí, México

*Corresponding author's e-mail address: arantzazu.luzuriaga@urjc.es

Associate Editor: Nishanta Rajakaruna

Populations & Communities. Chief Editor: Jean Burns

Abstract

Previous studies found that plant communities on infertile soils are relatively resistant to climatic variation due to stress tolerance adaptations. However, the species assemblies in gypsum soil habitats require further investigation. Thus, we considered the following questions. (1) Do harsher arid conditions determine the characteristics of the species that form plant assemblages? (2) Is the selection of the species that assemble in arid conditions mediated by their ability to grow on gypsum soils? (3) Is the selection of species that assemble in harsher conditions related to phylogenetically conserved functional traits? Perennial plant communities were analysed in 89 gypsum-soil sites along a 400 km climate gradient from the central to southeastern Iberian Peninsula. Each local assemblage was analysed in 30 × 30 m plots and described based on taxonomic, functional (soil plant affinity) and phylogenetic parameters. The mean maximum temperatures in the hottest month, mean annual precipitation and their interaction terms were used as surrogates for the aridity conditions in generalized linear models. In the hottest locations, the gypsophily range narrowed and the mean gypsophily increased at the community level, thereby suggesting the filtering of species and the dominance of soil specialists in the actual plant assemblages. Drier sites had higher taxonomic diversity. The species that formed the perennial communities were close in evolutionary terms at the two ends of the aridity gradient. The mean maximum temperatures in the hottest month had the main abiotic filtering effect on perennial plant communities, which was mediated by the ability of species to grow on gypsum soils, and thus gypsum specialists dominated the species assemblies in the hottest locations. In contrast, the perennial communities on gypsum soils were relatively resistant to changes in precipitation. Our findings suggest that the warmer environmental conditions predicted by global change models will favour gypsum specialists over generalists.

Keywords: Aridity gradient; assembly rules; community weighted mean (CWM); edaphic endemism; functional diversity; gypsum soil; Mediterranean; phylogenetic diversity; semiarid; soil affinity

Introduction

Ecological assembly rules represent the biotic and abiotic processes that prevent particular species from occurring in realized assemblages (*sensu* Keddy 1992; Wilson 1999). They are the ultimate expression of the existence of niche differences

and ecological filters that act hierarchically at different temporal and spatial scales (Gotzenberger *et al.* 2012; HilleRisLambers *et al.* 2012). Environmental filtering has been evaluated along fertility gradients (Spasojevic and Suding 2012; Gerhold *et al.* 2013; Price *et al.* 2014), but it has rarely been tested for particular

Received: 8 January 2020; Editorial decision: 28 April 2020; Accepted: 18 May 2020

© The Author(s) 2020. Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

restrictive soils such as serpentines (see [Fernandez-Going et al. 2012](#)) or gypsum soils (see [Luzuriaga et al. 2012; 2015](#), and [Peralta et al. 2019](#), for annual plants; [Delalandre and Montesinos-Navarro 2018](#), for plant–plant interactions). This is surprising because a high proportion of the taxonomic plant diversity in many hotspots depends on these restrictive soil habitats ([Médail and Quézel 1999](#); [Cacho and Strauss 2014](#)). Furthermore, the Intergovernmental Panel on Climate Change predicts that these special soil habitats will be greatly affected by intensified dryness ([IPCC 2014](#)), mostly because of their island-like structure ([Luzuriaga et al. 2018](#)). Gypsum outcrops are widespread restrictive soils throughout the world that mostly occur in arid and semiarid conditions ([Herrero et al. 2009](#)). Gypsum soils impose harsh abiotic conditions for plants (i.e. gypsophytes), and thus they provide an ideal model for analysing community assembly processes under special conditions ([Escudero et al. 2015](#)).

It is well known that edaphic specialists have stress-tolerant functional attributes ([Grime et al. 2000; 2008](#); [Escudero et al. 2015](#); [Sianta and Kay 2019](#)) and the assembly of plant species on gypsum soils depends greatly on the soil affinity or soil specialization characteristics of each species ([Palacio et al. 2007](#); [Luzuriaga et al. 2015](#) for gypsum soils; [Fernandez-Going et al. 2012](#) for serpentine plant communities). The mechanism that induces the restrictive occurrence of some species on gypsum soils is called gypsophily ([Merlo et al. 2009](#)). Thus, we can assume that gypsophily integrates the characteristics that a species needs to become established in these harsh environments (for more details, see [Escudero et al. 2015](#)). The gypsum dependence of species appears to be connected with evapotranspiration ([Escudero et al. 2015](#)), so in the present study, we hypothesized that the degree of specialization by the species in an assembly should vary with climate. In order to test this hypothesis and evaluate whether the assembly rules shift along climate gradients, we used the gypsum preference of each species as a plant functional trait. To determine the soil preferences of each species, we used the species-specific gypsophily values assigned by a committee of expert botanists who quantified the soil preferences ([Mota et al. 2011](#)). This so-called gypsophily index (GI) represents a species-specific plant functional property that is crucial for species establishment (including survival, growth and reproduction) ([Palacio et al. 2007](#)), and its analysis using similar techniques to other plant functional traits may help us to understand community assemblies on restrictive substrates (see [Luzuriaga et al. 2015](#)). We combined this GI with tools that were originally developed for describing trait diversity patterns among communities in order to identify the filters (i.e. assembly rules) that operate on perennial plant assemblages along an aridity gradient in the Iberian Peninsula. Xerophily seems to be a major component for integrating the gypsophily syndrome ([Escudero et al. 2015](#)), so we expected that the gypsophily range (GR) of co-occurring species would narrow and that the mean community gypsophily values would increase at the aridest end of our climate gradient.

Phylogenetic diversity is a surrogate for the evolutionary relationships among the species that form an actual community and it provides valuable information about the contribution of trait evolution to the community structure and composition ([Webb et al. 2002](#)). Community phylogenetic analysis alone may only provide limited insights into the mechanisms responsible for forming plant assemblages (see [Mayfield and Levine 2010](#); [Weiher et al. 2011](#)), but it may yield information regarding environmental filtering processes when combined with the interpretations of specific traits because phylogenetic diversity integrates the

patterns found in the traits of all species ([Kraft et al. 2007](#); [Kembel 2009](#)). It is assumed that closer species in evolutionary terms are usually more functionally similar compared with those that are more distantly related ([Connolly et al. 2011](#); [Gerhold et al. 2015](#); but see [Violle et al. 2011](#)). Consequently, according to the environmental filtering hypothesis ([Keddy 1992](#)), it is expected that the phylogenetic diversity will reduce if the environment becomes harsher because only species from certain clades may be adapted to these conditions.

The main aim of the present study was to determine how perennial plant communities on gypsum soils vary according to the climate conditions in terms of precipitation and the mean maximum temperatures in the hottest month. It is well established that both of these climate factors explain additional and complementary proportions of the variability in plant performance ([Munson et al. 2011, 2013](#)). Thus, we assessed the changes in taxonomic, functional and phylogenetic diversity along an aridity gradient in terms of the mean maximum temperatures and mean annual precipitation, which could also provide insights into the possible responses of these communities to ongoing climate change (see [Butterfield 2015](#)). The following questions were addressed in this study. (1) Do the harsher arid conditions (in terms of the mean maximum temperature in the hottest month and mean annual precipitation) in gypsum systems determine the characteristics of the species that form plant assemblages? (2) Is the selection of the species that assemble in more arid conditions mediated by their ability to grow on gypsum soils (i.e. soil affinity index)? (3) Is the selection of the species that assemble in harsher conditions related to phylogenetically conserved functional traits? Understanding the effects induced by climate on the plant community assemblages that inhabit restrictive soil types may be critical for predicting how this unique plant biota will respond to ongoing climate change.

Methods

Study area

We studied a climate gradient covering a geographical length of 400 km, which included all existing gypsum island regions from the centre to the southeastern tip of the Iberian Peninsula. In total, 89 sites were selected on soils derived from gypsum outcrops classified as Xeric Haplogypsid ([Soil Survey Staff 1994](#)), which were located on southerly oriented gentle slopes. In this study, the aridity gradient comprised the broadest possible range of arid conditions in the Iberian Peninsula compatible with gypsophilous vegetation ([Rivas-Martinez and Costa 1970](#); [Escudero et al. 2015](#)). The mean annual precipitation ranged from 195 to 565 mm and the mean maximum temperatures in the hottest month ranged from 30.8 to 34.5 °C ([www.aemet.com](#)). Aridity was defined as the joint effect of less annual precipitation together with higher mean maximum temperatures in the hottest month, which is more critical for plant development than the mean annual temperatures in terms of water availability in the soil. The perennial vegetation comprised gypsophilous scrubland dominated by *Thymus lacaitae*, *Helianthemum squamatum*, *Lepidium subulatum* and *Gypsophila struthium* in the centre of Spain, and by *Teucrium libanitis*, *Herniaria fruticosa* and *Fumana hispida* in the southeast [see [Supporting Information—Table S1](#)]. The vegetation structure in our study system was typical of gypsum outcrops and drylands, where it comprised a matrix of plant patches on large areas of bare soil ([Escudero et al. 2015](#)), with total cover values ranging between 11 and 52.2 %.

Sampling design

At each site, we established a square plot measuring 30 × 30 m within homogeneous vegetated areas. We assessed the composition and structure of perennial vascular plants using four 30-m long linear transects parallel to the slope and 8 m apart in each plot. For each transect, we recorded the intercept length of every perennial plant in contact with the transect line. Species-specific cover was estimated visually in 12 quadrats (1.5 × 1.5 m) placed in each plot, with three quadrats per transect line. The total plant cover in the whole plot was determined as the proportion of the four transects in contact with perennial shrubs. Total plant cover was included in the statistical models as an estimate of the productivity in each plot. Climate variables were estimated with the climate simulation model CLIMOEST (developed for the Iberian Peninsula) by inputting the altitude, geographic coordinates and hydrographic basin. This simulation model estimates climate parameters derived from monthly temperature and rainfall values over the latest 50 years (Sánchez-Palomares et al. 1999).

Taxonomic, functional and phylogenetic diversity indices

Each taxonomic, functional, and phylogenetic diversity parameter was calculated at the plot level (30 × 30 m). Perennial diversity was calculated using the inverse Simpson index as follows:

$$\text{Inverse Simpson} = \frac{1}{\sum_{i=1}^S p_i^2},$$

where S is the number of species at the plot level and p_i is the proportion of species i . The diversity index was computed using the *vegan* package (Oksanen et al. 2016) in R v.3.4.1 (R Development Core Team 2018).

In order to characterize the affinity of each species for gypsum soils, we used the GI for Iberian gypsophilous species proposed by Mota et al. (2011) [see Supporting Information—Table S1]. The GI ranges from 1 (species that avoid gypsum soils) to 5 (species strictly linked to gypsum substrates, which are also called gypsophytes). Species with GI values from 2 to 4 are called gypsovags, i.e. species that tolerate gypsum and that can be established effectively in these types of soils, but that usually occur on other substrate types as well. In our study, no species had a GI value of 1 because we actively selected gypsum soils for our sampling plots.

The affinity of species for gypsum soils was described at the community level using three parameters. In particular, we aimed to describe the range of soil affinity values in each species assembly (GR), the community mean gypsophily (CMG; weighted mean) and the gypsophily diversity (GD) in each actual assembly. These indices are not correlated with each other and they are independent of the species richness (Mason et al. 2005), and they allowed us to identify the main assembly processes related to the abundances of species (Mason et al. 2013). Mason et al. (2013) defined functional richness as the amount of a niche space filled by species in a realized assembly. Based on this definition, we computed the GR as analogous to the functional richness in order to quantify the amount of space used by each assemblage in the gypsophily gradient. We calculated the range of gypsophily values in each assembly as follows.

$$\text{GR} = \max(\text{GI}) - \min(\text{GI})$$

The GR values can range between 0 (all species in the assemblage have the same GI) and 3 (at least two species in the assemblage have extreme GI values).

The CMG index is analogous to the community weighted mean index (Lavorel et al. 2008), where it weights the GIs of species based on their relative abundance in the assemblage to obtain a single community level value. CMG was calculated as follows:

$$\text{CMG} = \sum_{i=1}^n (G_i \times p_i),$$

where p_i is the relative contribution of species i to the community and G_i is the GI of species i . CMG values can range between 2 (only species with sporadic presence on gypsum soils in the community) and 5 (only gypsophytes in the community).

The GD was employed to describe the divergence of the plant species cover along the gypsophily gradient in each assemblage. It was computed as the Rao quadratic diversity (Botta-Dukát 2005):

$$\text{GD} = \sum_{i,j}^S d_{ij} \times p_i \times p_j,$$

where S is the species richness and d_{ij} is the difference in the GI values between species i and j , which was calculated as follows.

$$d_{i,j} = (G_i - G_j)^2$$

These three community-level functional parameters were computed using the FD package (Laliberté et al. 2014) in R (R Development Core Team 2018).

We calculated the standardized effect size for GR and GD in order to standardize these values and avoid bias related to the species number in each plot. The GR and GD values observed were compared with those obtained after 10 000 randomizations of the species abundances (see Mason et al. 2013, for more details).

A phylogenetic tree was built for all 111 identified species in our study using the V.PhylMaker R package (Jin and Qian 2019). This statistical package attaches the species in a user-supplied list to a megatree of 74 533 vascular plant species where the branch lengths correspond to the evolutionary divergence time between branches. Three phylogenetic diversity indices were used: phylogenetic species variability (PSV), phylogenetic species richness (PSR) and net relatedness index (NRI). The PSV (Helmus et al. 2007) measures the phylogenetic relatedness of the species in a local assemblage, where the PSV equals 1 when all of the species in a sample are unrelated (i.e. a star phylogeny) and it approaches zero as species become more related in the phylogeny. PSR is the product of the PSV and species richness, and it can be considered as the species richness in a sample after discounting species relatedness. The PSR is maximized at the species richness of the sample and it decreases towards zero as the relatedness increases.

In general, the phylogenetic diversity of assemblages is correlated with the species richness, so we calculated the NRI values for our data. This index indicates whether the phylogenetic diversity in an assemblage is greater or less than expected regardless of the species richness (Webb 2000):

$$\text{NRI} = -1 \times \frac{\text{MPD} - \text{MPD}_{\text{rnd}}}{\text{sdMPD}_{\text{rnd}}},$$

where MPD_{rnd} are the means of the mean pairwise distance (MPD) values based on 999 randomly generated assemblages and the sdMPD_{rnd} are the standard deviations of the 999 MPDs obtained from those assemblages. Thus, negative NRI values indicate higher than expected phylogenetic diversity in an assemblage given the species richness of that assemblage.

The random assemblages generated in the null models were generated with the independent swap algorithm by drawing the same number of species from the pool as the number of species in the observed community, where the observed community occupancy rates were fixed (Gotelli and Entsminger 2001). These phylogenetic diversity indices were computed at the plot level using the Picante package (Kembel et al. 2010) in R (R Development Core Team 2018).

Statistical analysis

We constructed generalized linear models based on the taxonomic richness, inverse Simpson index, CMG, GR, GD, PSR, PSV and NRI indices. We used the mean maximum temperatures in the hottest month and the mean annual precipitation during the last 50 years, as well as the interaction terms between both variables, as surrogates for the aridity conditions in each plot. We checked for correlations between both explanatory variables in order to avoid any increase in the variance (Pearson's correlation coefficient: $R^2 = 0.17$, $t = 1.6$, and $P = 0.11$). Each model included the total plant cover in the plot as a surrogate of plant productivity in order to statistically control for the effects of productivity, thereby allowing us to evaluate the effects of temperature and precipitation regardless of the well-known productivity effects. The distribution error and link function with the best fits to our data were used for each generalized linear model (see Table 1).

Results

In total, 111 perennial plant species from 34 families were recorded along the aridity gradient on gypsum soils [see Supporting Information—Table S1]. Sixteen species preferred gypsum soils and they appeared rarely on other substrate types, whereas the remainder were capable of growing on substrates other than gypsum. The mean maximum temperature and annual precipitation did not affect the taxonomic diversity and PSR after controlling for productivity effects by including the total vegetation cover in each plot in our models. However, we detected significant effects of the mean maximum temperatures on the inverse Simpson index, and on the functional and phylogenetic diversity indices (Table 1; Fig. 1). High mean maximum temperatures were related to lower taxonomic

diversity and narrower ranges for the gypsophily indices. Thus, in the hottest environments, the GR narrowed at the community level. Furthermore, the CMG values showed that soil specialists (species with high gypsophily values) dominated the plant assemblages in hot conditions. In contrast, lower mean annual precipitation was associated with higher taxonomic diversity. Phylogenetic diversity was low at both ends of the gradient, at the most arid (high mean maximum temperatures and low precipitation) and at the least arid end (low mean maximum temperatures and high precipitation) (Fig. 1). Thus, the species that formed the actual perennial assemblages were closer in evolutionary terms at the most arid end of the gradient, and the species at the least arid end of the gradient were also phylogenetically closer to each other.

Discussion

Precipitation is usually considered a primary driver of the vegetation dynamics in semiarid grasslands (Chesson et al. 2004; Miranda et al. 2011; Luzuriaga et al. 2012; 2015; Mulhouse et al. 2017; Peralta et al. 2019). However, we found that it did not affect the species richness and functional and phylogenetic diversity after statistically controlling for productivity effects. Our results agree with those reported by Fernandez-Going et al. (2012) who observed that the community composition on low-fertility serpentine soils generally varied less in response to precipitation than communities on more fertile soils. In addition, our findings support the prediction by Grime et al. (2000, 2008) that plant communities on nutrient-poor soils may be relatively resistant to changes in precipitation. This ability could be explained by the development of stress-tolerant functional traits (Damschen et al. 2012; Peralta et al. 2019), but other factors such as strong nutrient limitation may reduce the potential impact of changes in rainfall on plant growth. Other studies found that communities responded to a change in one resource only after the removal of the limitation due to another resource or condition (Klanderud and Totland 2005; Going et al. 2009). Nutrient availability and precipitation are closely linked because low precipitation is known to reduce the availability of nutrients due to water limitation restricting soil microbial processes (de Dato et al. 2006; Sardans et al. 2008) and decreasing the mobility of nutrients. Thus, plant adaptations to infertile

Table 1. Chi-square values obtained by the generalized linear models for taxonomic, functional and phylogenetic diversity indices. Each model included the total plant cover in the plot as a covariable in order to statistically control for the differences in the vegetation productivity along the aridity gradient. T: Mean maximum temperatures in the hottest month; P: mean annual precipitation; GR: gypsophily range; CMG: community mean gypsophily index; GD: gypsophily diversity; PSR: phylogenetic species richness; PSV: phylogenetic species variability; NRI: net relatedness index. The error distributions (Family) and link functions assumed in the models are indicated. Id: identity link function; Log: logarithmic link function. The signs of the coefficients are shown in parentheses. * $0.01 < P < 0.05$; ** $0.001 < P < 0.01$; *** $P < 0.001$.

	Family (link)	T	P	T × P	Cover
Taxonomic indices					
Richness	Poisson (Log)	1.06	2.6	0.9	(+) 10.9***
Diversity	Gaussian (Id)	(-) 9.3**	(-) 5.1*	2.3	0.97
Functional indices					
GR	Gaussian (Id)	(-) 6.3*	0.9	1.1	0.12
CMG	Quasipoisson (Log)	(+) 11.1***	0.2	0.1	0.2
GD	Gaussian (Id)	2.1	0.1	3.2	
Phylogenetic indices					
PSR	Gaussian (Id)	2.2	0.4	2.3	(+) 5.3*
PSV	Gaussian (Id)	1.9	0.7	(+) 4.5*	0.09
NRI	Gaussian (Id)	2.4	0.2	(-) 7.3**	0.7




Figure 1. Graphic representation of the interaction between the mean maximum temperature in the hottest month and mean annual precipitation at the sampling locations based on: (A) the phylogenetic species variability (PSV) and (B) the Net Relatedness Index (NRI) in each plot. Circles represent the observed values. Lines join points with the same predicted PSV (A) or NRI (B) values based on our generalized linear models. Darker areas represent higher predicted values and lighter areas lower predicted values for PSV and NRI indices.

soils are probably closely related to adaptations to low water availability conditions (Chapin 1991).

The taxonomic diversity decreased as the water availability increased, which may have been the combined result of two processes acting simultaneously: (i) many gypsophilous species do not fully exploit the water availability in rainier sites because their stress-tolerant nature will constrain their ability to utilize this resource and they may be displaced by calcophylous plant species (Dalgleish and Hartnett 2006; Bai et al. 2008; Nippert and Holdo 2015); and (ii) the higher water availability may benefit competitive species that are particularly efficient at exploiting resources and occupying space, such as the perennial tussock *Macrochloa tenacissima*. Thus, water limitation may be crucial for the maintenance of high plant diversity on gypsum soils in these systems, as shown by Eskelinen and Harrison (2015) who suggested that water and nutrient colimitation is the mechanism responsible for plant community resistance to changes in precipitation. Some studies have shown that communities on infertile soils are more diverse (Cacho and Strauss 2014), which may confer resistance to climatic variability by enhancing the potential for complementarity among the responses of species (Tilman and Downing 1994; McCann 2000).

Our results provide good evidence for the filtering effect of the mean maximum temperatures on gypsum plant communities (see also Moles et al. 2014, Butterfield and Munson 2016), where the relative abundance of gypsum specialist plants increased towards the hottest locations, regardless of the mean annual precipitation. Temperatures can induce moisture stress in many plant species (De Boeck et al. 2007) and gypsum specialists are equipped with stress-tolerant attributes (e.g. naked buds, deeply rooted tap roots, succulence and persistent soil seed banks; see Palacio et al. 2007; Escudero et al. 2015; Peralta et al. 2016), so

heat may select the gypsophilous flora. Indeed, our results agree with those obtained in other studies of the distribution of strict gypsophiles (de Luis et al. 2019) and the regeneration stages of five gypsophiles (Sánchez et al. 2017). In general, gypsophily has been interpreted as an adaptive syndrome defined by traits that allow specialist plants to root in the hard physical crust (i.e. the physical hypothesis; Meyer 1986; Romão and Escudero 2005) and to cope with the ion imbalance that characterizes gypsisoils (i.e. the chemical hypothesis; Duvigneaud and Denaeyer-De Smet 1968; Boukhris and Lossaint 1975; Palacio et al. 2007). However, the participation of climate factors themselves in the modulation of gypsophily has received little attention, probably because it is broadly assumed that the gypsophilous flora is well adapted to stressful climate conditions. Future investigations should elucidate the effect of temperature on the modulation of gypsophily as an independent driver or as an environmental component that exacerbates other factors.

It should also be noted that the GD was not affected by the mean maximum temperature or annual precipitation, thereby suggesting that the actual local assemblages contained species that differed in terms of their tolerance of gypsum. Facilitation under the canopy of shrubs may have led to the presence of gypsovags beneath their canopies (Romão and Escudero 2005; Luzuriaga et al. 2015). Furthermore, the GD values were stable even though there were remarkable shifts in the species composition along the aridity gradient (see Rivas-Martínez and Costa 1970), which was probably related to the divergent biogeographical history of the gypsum geological outcrops in the Iberian Peninsula (Herrero et al. 2009).

In this study, we found that evolutionarily close relatives coexisted in perennial plant communities (low phylogenetic diversity) at the most arid end of the aridity gradient (high mean maximum temperatures and low annual precipitation), and a similar situation occurred in the least arid locations (low mean maximum temperatures and high mean precipitation). These results were probably due to the intensification of different natural environmental filters operating at both extremes of the gradient. Thus, our results suggest that phylogenetic clustering might not have been induced by abiotic filtering under harsh conditions, but instead it could have been due to biotic interactions in more productive conditions. Bernard-Verdier et al. (2012) found similar trends in a Mediterranean rangeland. The coexistence of close relatives in resource-poor sites suggests that community membership is limited by a requirement for shared stress tolerance traits, i.e. habitat phylogenetic filtering (Grime et al. 2008; Butterfield et al. 2013). However, at the least arid end of the climate gradient, competition may have been a strong biotic filter, thereby leading to phylogenetic convergence via mechanisms for equalizing fitness (Chesson 2000; Grime 2006). Abiotic or biotic filtering may have relaxed as the aridity gradient changed to an intermediate level and this could have allowed the phylogenetic structure of the community to expand. According to the results obtained in the present study, the mean maximum temperatures had the main abiotic filtering effect on perennial plant communities, which was mediated by the ability of species to grow on gypsum soils (i.e. soil affinity index); thus, gypsum specialists dominated the species assemblages in the hottest locations. In contrast, the perennial communities on gypsum soils may have been relatively resistant to changes in precipitation. Therefore, our findings suggest that the warmer environmental conditions predicted by global change models may favour gypsum specialists (see de Luis et al. (2019) for an example of a strict gypsophile), thereby making the plant communities more gypsophilous in semiarid systems. However, several processes may interact, such as climate variability

(Adler et al. 2006), extreme climatic events (Evans et al. 2011), nutrient availability (Delgado-Baquerizo et al. 2013), the soil microbial community (Zogg et al. 1997; Castro et al. 2010) and the trade-off between survival and plant growth (Lloret et al. 2012; Benavides et al. 2015), and thus the possible consequences for perennial plant communities are difficult to predict. In conclusion, the high taxonomic diversity of Mediterranean gypsum systems appears to be related to stressful arid conditions, and the tolerance of species for these conditions may be summarized using the GI for each species.

Data Availability

Data used in this article are available at the URJC public repository under the following link: http://repositories.biodiversos.org/Luzuriaga_A.L/

Supporting Information

The following supporting information is available in the online version of the article—

Table S1. Perennial plant species that occurred in the 89 plots located on gypsum soils and their gypsophily indices (GI).

Sources of Funding

This study was supported by the Ministerio de Economía y Competitividad AGORA project (CGL2016-77417-P—Spanish Government), REMEDINAL project (TE-CM S2018/EMT-4338—Madrid Regional Government), Ministerio de Ciencia e Innovación PHENOTYPES project (PGC2018-099115-B-I00-P—Spanish Government) and GYPWORLD project (H2020-MSCA-RISE-2017-777803—European Commission).

Acknowledgements

We thank Esmeralda Martínez-Duro and Emilia Alquézar for field assistance and Mario Blanco for obtaining the climate variables with CLIMOEST. Marcelino de la Cruz for assistance with making isocline figures with R. We thank the Spanish Meteorological Agency (AEMET) for providing climatic data.

Contributions by the Authors

A.E.: conceived the main ideas. A.L.L., J.F. and P.F.: conducted field sampling. A.L.L.: performed the statistical analysis and led the writing of the manuscript. All co-authors contributed to the main text.

Conflict of Interest

None declared.

Literature Cited

Adler PB, HilleRisLambers J, Kyriakidis PC, Guan Q, Levine JM. 2006. Climate variability has a stabilizing effect on the coexistence of prairie grasses *Proceedings of The National Academy of Sciences of the United States of America* **103**:12793–12798.

Bai Y, Wu J, Xing Q, Pan Q, Huang J, Yang D, Han X. 2008. Primary production and rain use efficiency across a precipitation gradient on the Mongolia Plateau. *Ecology* **89**:2140–2153.

Benavides R, Escudero A, Coll L, Ferrandis P, Gouriveau F, Hódar JA, Ogaya R, Rabasa SG, Granda E, Santamaría BP, Martínez-Vilalta J, Zamora R, Espelta JM, Peñuelas J, Valladares F. 2015. Survival vs. growth trade-off in early recruitment challenges global warming impacts on Mediterranean mountain trees. *Perspectives in Plant Ecology, Evolution and Systematics* **17**:369–378.

Bernard-Verdier M, Navas ML, Vellend M, Violle C, Fayolle A, Garnier E. 2012. Community assembly along soil depth gradient: contrasting patterns of plant trait convergence and divergence in a Mediterranean rangeland. *Journal of Ecology* **100**:1422–1433.

Botta-Dukát Z. 2005. Rao's quadratic entropy as a measure of functional diversity based on multiple traits. *Journal of Vegetation Science* **16**:533–540.

Boukhris M, Lossaint P. 1975. Aspects ecologiques de la nutrition minerale de plantes gypsicoles di Tunisie. *Revue d'Ecologie et de Biologie du Sol* **12**:329–348.

Butterfield BJ. 2015. Environmental filtering increases in intensity at both ends of climatic gradients, though driven by different factors, across woody vegetation types of the southwest USA. *Oikos* **10**:1374–1382.

Butterfield BJ, Cavieres LA, Callaway RM, Cook BJ, Kikvidze Z, Lortie CJ, Michalet R, Pugnaire FI, Schöb C, Xiao S, Zaitchek B, Anthelme F, Björk RG, Dickinson K, Gavilán R, Kanka R, Maalouf JP, Noroozi J, Parajuli R, Phoenix GK, Reid A, Ridenour W, Rixen C, Wipf S, Zhao L, Brooker RW. 2013. Alpine cushion plants inhibit the loss of phylogenetic diversity in severe environments. *Ecology Letters* **16**:478–486.

Butterfield BJ, Munson SM. 2016. Temperature is better than precipitation as a predictor of plant community assembly across a dryland region. *Journal of Vegetation Science* **27**:938–947.

Cacho NI, Strauss SY. 2014. Occupation of bare habitats, an evolutionary precursor to soil specialization in plants. *Proceedings of the National Academy of Sciences of the United States of America* **111**:15132–15137.

Cadotte MW, Cavender-Bares J, Tilman D, Oakley TH. 2009. Using phylogenetic, functional and trait diversity to understand patterns of plant community productivity. *PLoS One* **4**:e5695.

Castro HF, Classen AT, Austin EE, Norby RJ, Schadt CW. 2010. Soil microbial community responses to multiple experimental climate change drivers. *Applied and Environmental Microbiology* **76**:999–1007.

Chapin III FS. 1991. Integrated responses of plants to stress. *Bioscience* **41**:29–36.

Chesson P. 2000. Mechanisms of maintenance of species diversity. *Annual Review Ecology and Systematics* **31**:343–366.

Chesson P, Gebauer RL, Schwinning S, Huntly N, Wiegand K, Ernest MS, Sher A, Novoplansky A, Weltzin JF. 2004. Resource pulses, species interactions, and diversity maintenance in arid and semi-arid environments. *Oecologia* **141**:236–253.

Connolly J, Cadotte MW, Brophy C, Dooley A, Finn J, Kirwan L, Roscher C, Weigelt A. 2011. Phylogenetically diverse grasslands are associated with pairwise interspecific processes that increase biomass. *Ecology* **92**:1385–1392.

Dalgleish HJ, Hartnett DC. 2006. Below-ground bud banks increase along a precipitation gradient of the North American Great Plains: a test of the meristem limitation hypothesis. *The New Phytologist* **171**:81–89.

Damschen EI, Harrison S, Ackerly DD, Fernandez-Goñi BM, Anacker BL. 2012. Endemic plant communities on special soils: early victims or hardy survivors of climate change? *Journal of Ecology* **100**:1122–1130.

De Boeck H, Lemmens CMHM, Gielen B, Nijs I. 2007. Combined effects of climate warming and plant diversity loss on above- and below-ground grassland productivity. *Environmental and Experimental Botany* **60**:95–104.

De Dato G, Pellizzaro G, Cesaraccio C, Sirca C, De Angelis P, Duce P, Spano D, Scarascia-Mugnozza G. 2006. Effects of warmer and drier climate conditions on plant composition and biomass production in a Mediterranean shrubland community. *Forest Ecology* **3**:511–526.

de Luis M, Álvarez-Jiménez J, Martínez Labarga JM, Bartolomé C. 2019. Four climate change scenarios for *Gypsophila bermejoi* G. López (Caryophyllaceae) to address whether bioclimatic and soil suitability will overlap in the future. *PLoS One* **14**:e0218160.

Delalandre L, Montesinos-Navarro A. 2018. Can co-occurrence networks predict plant-plant interactions in a semi-arid gypsum community? *Perspectives in Plant Ecology, Evolution and Systematics* **31**:36–43.

- Delgado-Baquerizo M, Maestre FT, Gallardo A, Bowker MA, Wallenstein MD, Quero JL, Ochoa V, Gozalo B, García-Gómez M, Soliveres S, García-Palacios P, Berdugo M, Valencia E, Escolar C, Arredondo T, Barraza-Zepeda C, Bran D, Carreira JA, Chaieb M, Conceição AA, Derak M, Eldridge DJ, Escudero A, Espinosa CI, Gaitán J, Gatica MG, Gómez-González S, Guzman E, Gutiérrez JR, Florentino A, Hepper E, Hernández RM, Huber-Sannwald E, Jankju M, Liu J, Mau RL, Miriti M, Moneris J, Naseri K, Noumi Z, Polo V, Prina A, Pucheta E, Ramírez E, Ramírez-Collantes DA, Romão R, Tighe M, Torres D, Torres-Díaz C, Ungar ED, Val J, Wamiti W, Wang D, Zaady E. 2013. Decoupling of soil nutrient cycles as a function of aridity in global drylands. *Nature* 502:672–676.
- Duvigneaud P, Denaeyer-De Smet S. 1968. Essai de classification chimique (elements minéraux) des plantes gypsicolas du Bassin de l'Ebre. *Bulletin de la Societ e Royale de Botanique de Belgique* 101:279–291.
- Escudero A, Palacio S, Maestre FT, Luzuriaga AL. 2015. Plant life on gypsum: a review of its multiple facets. *Biological Reviews of the Cambridge Philosophical Society* 90:1–18.
- Eskelinen A, Harrison SP. 2015. Resource colimitation governs plant community responses to altered precipitation. *Proceedings of the National Academy of Sciences of the United States of America* 112:13009–13014.
- Evans SE, Byrne KM, Lauenroth WK, Burke IC. 2011. Defining the limit to resistance in a drought-tolerant grassland: long-term severe drought significantly reduces the dominant species and increases ruderals. *Journal of Ecology* 99:1500–1507.
- Fernandez-Going BM, Anacker BL, Harrison SP. 2012. Temporal variability in California grasslands: soil type and species functional traits mediate response to precipitation. *Ecology* 93:2104–2114.
- Gerhold P, Cahill JF, Winter M, Bartish I, Prinzing A. 2015. Phylogenetic patterns are not proxies of community assembly mechanisms (they are far better) *Functional Ecology* 29:600–614.
- Gerhold P, Price JN, Püssa K, Kalamees R, Aher K, Kaasik A. 2013. Functional and phylogenetic community assembly linked to changes in species diversity in a long-term resource manipulation experiment *Journal of Vegetation Science* 24:43–852.
- Going BM, Hillerislambers J, Levine JM. 2009. Abiotic and biotic resistance to grass invasion in serpentine annual plant communities. *Oecologia* 159:839–847.
- Gotelli NJ, Entsminger GL. 2001. Swap and fill algorithms in null model analysis: rethinking the knight's tour. *Oecologia* 129:281–291.
- Götzenberger L, de Bello F, Brathen KA, Davison J, Dubuis A, Guisan A, Leps J, Lindborg R, Moora M, Zobel M. 2012. Ecological assembly rules in plant communities approaches, patterns and prospects. *Biological Reviews* 87:111–127.
- Grime JP. 2006. Trait convergence and trait divergence in the plant community: mechanisms and consequences. *Journal of Vegetation Science* 17:255–260.
- Grime JP, Brown VK, Thompson K, Masters GJ, Hillier SH, Clarke IP, Askew AP, Corker D, Kieley JP. 2000. The response of two contrasting limestone grasslands to simulated climate change. *Science (New York, N.Y.)* 289:762–765.
- Grime JP, Fridley JD, Askew AP, Thompson K, Hodgson JG, Bennett CR. 2008. Long-term resistance to simulated climate change in an infertile grassland. *Proceedings of the National Academy of Sciences of the United States of America* 105:10028–10032.
- Helmus MR, Bland TJ, Williams CK, Ives AR. 2007. Phylogenetic measures of biodiversity. *American Naturalist* 169:E68–E83.
- Herrero J, Artieda O, Hudnall WH. 2009. Gypsum, a tricky material. *Soil Science Society of America Journal* 73:1757–1763.
- HilleRisLambers J, Adler PB, Harpole WS, Levine JM, Mayfields MM. 2012. Rethinking community assembly through the lens of coexistence theory. *Annual Reviews of Ecology, Evolution and Systematics* 43:227–248.
- IPCC 2014. *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, RK Pachauri, LA Meyer, eds.]. Geneva, Switzerland: IPCC, 151 pp.
- Jin Y, Qian H. 2019. V. PhyloMaker: an R package that can generate very large phylogenies for vascular plants. *Ecography* 42:1353–1359.
- Keddy PA. 1992. Assembly and response rules: two goals for predictive community ecology. *Journal of Vegetation Science* 3:157–164.
- Kemmel SW. 2009. Disentangling niche and neutral influences on community assembly: assessing the performance of community phylogenetic structure tests. *Ecology Letters* 12:949–960.
- Kemmel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, Blomberg SP, Webb CO. 2010. Picante: R tools for integrating phylogenies and ecology. *Bioinformatics (Oxford, England)* 26:1463–1464.
- Klanderud K, Totland Ø. 2005. Simulated climate change altered dominance hierarchies and diversity of an alpine biodiversity hotspot. *Ecology* 86:2047–2054.
- Kraft NJ, Cornwell WK, Webb CO, Ackerly DD. 2007. Trait evolution, community assembly, and the phylogenetic structure of ecological communities. *The American Naturalist* 170:271–283.
- Lalibert e E, Legendre P, Shipley B. 2014. FD: measuring functional diversity from multiple traits, and other tools for functional ecology. R package version 1.0–12.
- Lavorel S, Grigulis K, McIntyre S, Williams NSG, Garden D, Dorrough J, Berman S, Quetier F, Thebault A, Bonis A. 2008. Assessing functional diversity in the field—methodology matters! *Functional Ecology* 22:134–147.
- Lloret F, Escudero A, Iriondo JM, Martínez-Vilalta J, Valladares F. 2012. Extreme climatic events and vegetation: the role of stabilizing processes. *Global Change Biology* 18:797–805.
- Luzuriaga AL, González JM, Escudero A. 2015. Annual plant community assembly in edaphically heterogeneous environments. *Journal of Vegetation Science* 26:866–875.
- Luzuriaga AL, Sánchez AM, López-Angulo J, Escudero A. 2018. Habitat fragmentation determines diversity of annual plant communities at landscape and fine spatial scales. *Basic and Applied Ecology* 29:12–19.
- Luzuriaga AL, Sánchez AM, Maestre FT, Escudero A. 2012. Assemblage of a semi-arid annual plant community, abiotic and biotic filters act hierarchically. *PLoS One* 7:e41270.
- Mason NWH, de Bello F, Mouillot D, Pavoine S, Dray S. 2013. A guide for using functional diversity indices to reveal changes in assembly processes along ecological gradients. *Journal of Vegetation Science* 24:794–806.
- Mason NWH, Mouillot D, Lee WG, Wilson JB. 2005. Functional richness, functional evenness and functional divergence: the primary components of functional diversity. *Oikos* 111:112–118.
- Mayfield MM, Levine JM. 2010. Opposing effects of competitive exclusion on the phylogenetic structure of communities. *Ecology Letters* 13:1085–1093.
- McCann KS. 2000. The diversity-stability debate. *Nature* 405:228–233.
- Médail F, Qu ezel P. 1999. Biodiversity hotspots in the Mediterranean Basin: setting global conservation priorities. *Conservation Biology* 13:1510–1513.
- Merlo ME, De Carrasco CG, Sola AJ, Jiménez-Sánchez ML, Rodríguez ML, Mota JF. 2009. Can gypsophytes distinguish different types of gypsum habitats? *Acta Botanica Gallica* 156:63–78.
- Meyer SE. 1986. The ecology of gypsophile endemism in the eastern Mojave Desert. *Ecology* 67, 1303–1313.
- Miranda JD, Armas C, Padilla FM, Pugnaire FI. 2011. Climatic change and rainfall patterns: effects on semi-arid plant communities of the Iberian Southeast. *Journal of Arid Environments* 75:1302–1309.
- Moles AT, Perkins SE, Laffan WS, Flores-Moreno H, Awasthy M, Tindall ML, Sack L, Pitman A, Kattge J, Bonser SP. 2014. Which is a better predictor of plant traits: temperature or precipitation? *Journal of Vegetation Science* 25:1167–1180.
- Mota JF, Sánchez-Gómez P, Guirado JS. 2011. *Diversidad vegetal de las yeseras ib ericas*. Almer a: ADIF-Mediterr aneo Asesores Consultores.
- Mulhouse JM, Hallett LM, Collins SL. 2017. The influence of seasonal precipitation and grass competition on 20 years of forb dynamics in northern Chihuahuan Desert grassland. *Journal of Vegetation Science* 28:250–259.
- Munson SM, Belnap J, Schelz CD, Moran M, Carolin TW. 2011. On the brink of change: plant responses to climate on the Colorado Plateau. *Ecosphere* 2:art68.
- Munson SM, Muldavin EH, Belnap J, Peters DP, Anderson JP, Reiser MH, Gallo K, Melgoza-Castillo A, Herrick JE, Christiansen TA. 2013. Regional signatures of plant response to drought and elevated temperature across a desert ecosystem. *Ecology* 94:2030–2041.
- Nippert JB, Holdo RM. 2015. Challenging the maximum rooting depth paradigm in grasslands and savannas. *Functional Ecology* 29:739–745.

- Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlenn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H. 2016. *Vegan: Community Ecology Package. Ordination methods, diversity analysis and other functions for community and vegetation ecologists*. Version 2.3–5.
- Palacio S, Escudero A, Montserrat-Martí G, Maestro M, Milla R, Albert MJ. 2007. Plants living on gypsum: beyond the specialist model. *Annals of Botany* **99**:333–343.
- Peralta AM, Sánchez AM, Luzuriaga AL, Escudero A. 2016. Factors driving species assemblage in Mediterranean soil seed banks: from the large to the fine scale. *Annals of Botany* **117**:1221–1228.
- Peralta AML, Sánchez AM, Luzuriaga AL, de Bello F, Escudero A. 2019. Evidence of functional species sorting by rainfall and biotic interactions: a community monolith experimental approach. *Journal of Ecology* **107**:2772–2788.
- Price JN, Gazol A, Tamme R, Hiiesalu I, Pärtel M. 2014. The functional assembly of experimental grasslands in relation to fertility and resource heterogeneity. *Functional Ecology* **28**:509–519.
- R Core Team. 2018. *A language and environment for statistical computing*. Vienna, AT: R Foundation for Statistical Computing.
- Rivas-Martínez S, Costa M. 1970. Comunidades gipsícolas del centro de España. *Anales del Instituto Botánico A.J. Cavanilles* **27**:193–224.
- Romao R, Escudero A. 2005. Gypsum physical soil surface crusts and the existence of gypsophytes in semi-arid central Spain. *Plant Ecology* **181**:1–11.
- Sánchez AM, Alonso-Valiente P, Albert MJ, Escudero A. 2017. How might edaphic specialists in gypsum islands respond to climate change? Reciprocal sowing experiment to infer local adaptation and phenotypic plasticity. *Annals of Botany* **120**:135–146.
- Sánchez-Palomares O, Sánchez-Serrano F, Carretero-Carrero MP. 1999. *Modelos y cartografía de estimaciones climáticas termoplumiométricas para la España peninsular*. INIA, Madrid, Spain: Colección Fuera de Serie.
- Sardans J, Peñuelas J, Estiarte M, Prieto P. 2008. Warming and drought alter C and N concentration, allocation and accumulation in a Mediterranean shrubland. *Global Change Biology* **14**:2304–2316.
- Sianta SA, Kay KM. 2019. Adaptation and divergence in edaphic specialists and generalists: serpentine soil endemics in the California flora occur in barer serpentine habitats with lower soil calcium levels than serpentine tolerators. *American Journal of Botany* **106**:690–703.
- Soil Survey Staff. 1994. *Keys to soil taxonomy*, 6th edn. Blacksburg: Pocahontas Press.
- Spasojevic MJ, Suding KN. 2012. Inferring community assembly mechanisms from functional diversity patterns: the importance of multiple assembly processes. *Journal of Ecology* **100**:652–661.
- Tilman D, Downing JA. 1994. Biodiversity and sustainability in grasslands. *Nature* **367**:363–365.
- Violle C, Nemergut DR, Pu Z, Jiang L. 2011. Phylogenetic limiting similarity and competitive exclusion *Ecology Letters* **14**:782–787.
- Webb CO. 2000. Exploring the phylogenetic structure of ecological communities: an example for rain forest trees. *The American Naturalist* **156**:145–155.
- Webb CO, Ackerly DD, McPeck MA, Donoghue MJ. 2002. Phylogenies and community ecology. *Annual Review of Ecology and Systematics* **33**:475–505.
- Weiher E, Freund D, Bunton T, Stefanski A, Lee T, Bentivenga S. 2011. Advances, challenges, and a developing synthesis of ecological community assembly theory. *Philosophical Transactions of the Royal Society B: Biological Sciences* **366**:2403–2413.
- Wilson JB. 1999. Assembly rules in plant communities. In: Weiher PA, Keddy E, eds. *Ecological assembly rules: perspectives, advances, retreats*. Cambridge, UK: Cambridge University Press, 130–164.
- Zogg GP, Zak DR, Ringelberg DB, MacDonald NW, Pregitzer KS, White DC. 1997. Compositional and functional shifts in microbial communities due to soil warming. *Soil Science Society of America Journal* **61**:475–481.