



**INSTITUTO POTOSINO DE INVESTIGACIÓN
CIENTÍFICA Y TECNOLÓGICA, A.C.**

POSGRADO EN CIENCIAS EN BIOLOGIA MOLECULAR

**Bacterial biodegradation of diisononyl phthalate:
molecular characterization and optimization**

Tesis que presenta

Marco Antonio Pereyra Camacho

Para obtener el grado de

Doctor en Ciencias en Biología Molecular

Director de la Tesis:

Dr. Antonio De León Rodríguez

San Luis Potosí, S.L.P., mayo de 2021



Constancia de aprobación de la tesis

La tesis “**Bacterial biodegradation of diisononyl phthalate: molecular characterization and optimization**” presentada para obtener el Grado de Doctor(a) en Ciencias en Biología Molecular fue elaborada **por M.C. Marco Antonio Pereyra Camacho** y aprobada el **17 de mayo de 2021** por los suscritos, designados por el Colegio de Profesores de la División de Biología Molecular del Instituto Potosino de Investigación Científica y Tecnológica, A.C.

Dr. Antonio De León Rodríguez
Director de la tesis

Dr. Victor Emmanuel Balderas Hernández
Miembro del Comité Tutorial

Dra. Sonia Lorena Arriaga García
Miembro del Comité Tutorial

Dr. Braulio Gutiérrez Medina
Miembro del Comité Tutorial



Créditos Institucionales

Esta tesis fue elaborada en el Laboratorio de Bioingeniería y Biotecnología Molecular de la División de Biología Molecular del Instituto Potosino de Investigación Científica y Tecnológica, A.C., bajo la dirección del Dr. Antonio De León Rodríguez. Se contó con el financiamiento del proyecto CONACyT Problemas Nacionales 4601.

Durante la realización del trabajo el autor recibió una beca académica del Consejo Nacional de Ciencia y Tecnología (No. 331852) y del Instituto Potosino de Investigación Científica y Tecnológica, A. C.

Dedications

I dedicate my thesis work to my family and friends. Especially to my parents and grandparents, for being my driving force and source of inspiration. To my dear siblings, Marialeidy and Claudio.

I also dedicate this thesis to my friends, classmates and colleagues who supported me during all this time.

I dedicate this work and give special thanks to my grandfather Rubén Alfonso Camacho Del Rincón, for being an example of life, my inspiration and for his words of encouragement. I carry you in my heart always grandfather.

To my great mentors and friends Antonio De Leon and Victor Balderas.

Thank you for everything!

R.I.P. RACR[†]♥ & CPS[†]♥

Acknowledgment

Thanks to Consejo Nacional de Ciencia y Tecnología (CONACYT) and Instituto Potosino de Investigación Científica y Tecnológica, A.C. (IPICYT). Thanks to the partial financial support from CONACYT Problemas Nacionales Grant 4601

I would like to thank Dr. Antonio De León Rodríguez for allowing me to be part of his research team, his teachings, patience and for his support during my studies. I wish to thank to the members of my thesis committee, Dr. Victor E. Balderas Hernández, Dra. Sonia Arriaga, Dr. Braulio Gutiérrez Medina, for their guidance, support, and contribution to this research.

Thanks to my family at the Bioengineering and Molecular Biotechnology Lab for their teachings, support and help all these years. Thanks to my classmates and colleagues for their valuable advice and for the good times we had together.

In addition, I would like to thank to my partners and friends of LabGeek for their support, complicity, unconditional friendship and all the adventures we had together. Also, thanks to my Rockstars team for their advice, support, and words of encouragement in this stage of my life.

Finally, I am extremely grateful to my family for their love, unconditional support, motivation, and encouragement during all this time.

Thank you for everything!

Content

	Page
Proof of approval of the thesis	ii
Institutional credits	iii
Exam certificate	iv
Dedications	v
Acknowledgment	vi
List of tables	viii
List of figures	ix
Annexes	x
Resumen	xi
Abstract	xii
Graphical abstract	xiii
1 Introduction	1
2. Hypothesis and aims	3
2.1 Hypothesis	3
2.2 Aims	3
2.2.1 General	3
2.2.2 Specific	3
3 Material and methods	4
3.1 Chemicals	4
3.2 Bacterial collection and formulation of DINP-degrading SSB-consortium	4
3.3 Enrichment culture of the SSB-consortium	5
3.4 Taxonomic identification of the SSB-consortium (16S Illumina sequencing)	5
3.5 Optimization of DINP biodegradation conditions	6
3.6 Biodegradation kinetics of DINP by the SSB-consortium	8
3.7 Analytical methods	9
4 Results	10
4.1 Identification and characterization of the SSB-consortium.	10
4.2 Optimization of operational conditions for DINP biodegradation	11
4.3 Biodegradation kinetics of DINP	17
4.4 Identification of DINP degradation intermediates and metabolic pathways	19
5 Discussion	22
6 Conclusion	25
7 References	25
8 Concluding remarks	35
9 Perspectives	36
10 Annexes	37

List of Tables

	Page
Table 1. Bacterial diversity of the SSB-consortium sample for NCBI BioProject PRJNA691434.	6
Table 2. Levels of independent variables used in the CCD for the optimization of DINP degradation.	7
Table 3. Matrix of the Composite Central Design and the response: % degradation of DINP.	13
Table 4. ANOVA of the % of DINP degradation obtained by Central Composite Design.	16
Table 5. Kinetic equations of DINP biodegradation at different initial DINP concentrations.	19
Table 6. Identification of DINP biodegradation metabolites by GC/MS.	20

List of Figures

	Page
Figure 1. Identification of the bacterial diversity of the SSB-consortium. (a) at family level; (b) at genus level.	11
Figure 2. Contour and 3D response surface plots showing the influence of pH, temperature, and initial DINP concentration on the % of DINP biodegradation by the SSB-consortium.	15
Figure 3. Growth of the SSB-consortium and DINP biodegradation kinetics started at 500 mg L ⁻¹ DINP: a) Evaluation of cell growth in MSM with or without DINP as the only carbon source, and b) degradation of DINP by the SSB-consortium under optimal conditions (31 °C, pH 7 and 500 mg L ⁻¹) in MSM. Non-inoculated MSM was employed as control (black squares). Error bars indicate the standard error of the three replicates.	18
Figure 4. Biodegradation of DINP at different initial concentrations performed by the SSB-consortium in MSM at 31 °C and pH 7. DINP concentrations used were: ▲ 250 mg L ⁻¹ , ● 500 mg L ⁻¹ and ○ 750 mg L ⁻¹ .	19
Figure 5. GC/MS chromatograms of DINP biodegradation by the SSB-consortium: a) 0 h, b) 72 h, c) 120 h, and d) 168 h of bacterial growth.	21
Figure 6. Proposed pathways for DINP degradation by the SSB-consortium. The dotted box indicates that the metabolite was not detected in this study.	22

Annexes

Scientific articles

1. Biodegradation of diisononyl phthalate by a consortium of saline soil bacteria: optimisation and kinetic characterization **37**
2. Biodegradation of recalcitrant compounds and phthalates by culturable bacteria isolated from *Liometopum apiculatum* microbiota **37**

Congress conferences

3. Bacterial biodegradation of endocrine disruptors di-(2-ethylhexyl) phthalate and di-isononyl phthalate. XXXII National Congress of Biochemistry (2018). Ixtapa Zihuatanejo, Gro. Mexico. **38**

Science communication conferences

4. Biodegradation - Phthalates in Agricultural Soils. III International Congress on Agroindustry: Industrial Trends, Biotechnology and Entrepreneurship (2020). Labens. Riobamba, Ecuador. **39**
5. Bioremediation 3.0. The Revolution of Systems Biology. 9th Edition of the International Biotechnology Congress QUORUM (2020). Instituto Tecnológico y de Estudios Superiores de Monterrey, Campus Querétaro. Querétaro, Mexico. **40**
6. Phthalates: a silent threat. 7th Biotechnology Week of Sonora Institute of Technology (2019). Cd. Obregon, Mexico. **41**

Resumen

Biodegradación bacteriana de diisononil ftalato: caracterización molecular y optimización

El diisononilftalato (DINP) es uno de los plastificantes más empleados en la producción de materiales plásticos y pertenece a los contaminantes ambientales más importantes. En este trabajo se presenta un consorcio de bacterias de suelos salinos (SSB) capaz de degradar el DINP. Los géneros del consorcio fueron *Serratia* sp., *Methylobacillus* sp., *Achromobacter* sp., *Pseudomonas* sp., *Stenotrophomonas* sp., *Methyloversatilis* sp., *Delftia* sp. y *Brevundimonas* sp. Se empleó la estrategia de metodología de superficie de respuesta (RSM) para optimizar y evaluar las condiciones de cultivo para la biodegradación del DINP. Las condiciones óptimas fueron un pH de 7.0, 31°C y una concentración inicial de DINP de 500 mg L⁻¹, lo que dio lugar a una biodegradación casi completa (99%) en 168 h. La degradación del DINP siguió un modelo cinético de primer orden, y la vida media fue de 12.76 h. Durante la biodegradación del DINP, se identificaron 4 compuestos derivados: monoisononilftalato, metilnonilftalato, iso-nanol y dimetil ftalato. El perfil de metabolitos indicó que el DINP se degradó a través de vías simultáneas de desesterificación y β -oxidación. Los resultados sugieren que el consorcio SSB podría ser útil para la biodegradación eficiente de los ambientes contaminados con DINP.

Palabras clave: Biodegradación; Consorcio; Degradación; Disruptores endocrinos; Plastificantes; Ftalatos.


Abstract

Bacterial biodegradation of diisononyl phthalate: molecular characterization and optimization

Di-isononyl phthalate (DINP) is one of plasticizers most employed in the production of plastic materials and belongs to the most important environmental contaminants. In this work, a consortium of saline soil bacterial (SSB) capable of degrading DINP is presented. The genera of SSB-consortium were *Serratia* sp., *Methylobacillus* sp., *Achromobacter* sp., *Pseudomonas* sp., *Stenotrophomonas* sp., *Methyloversatilis* sp., *Delftia* sp. and *Brevundimonas* sp. Response surface methodology (RSM) study was employed to optimise and evaluate the culture conditions to improve the biodegradation of DINP. The optimal conditions were a pH 7.0, 31°C and an initial DINP concentration of 500 mg L⁻¹, resulting in almost complete biodegradation (99%) in 168 h. DINP degradation followed a first-order kinetic model, and the half-life was 12.76 h. During the biodegradation of DINP, 4 derived-compounds were identified: monoisononyl phthalate, methyl nonyl phthalate, iso-nonanol, and dimethyl phthalate. The metabolite profiling indicated that DINP was degraded through simultaneous pathways of de-esterification and β -oxidation. Results suggest that the SSB-consortium could be useful for efficient biodegradation of the DINP-contaminated environments.

Keywords: Biodegradation; Consortium; Degradation; Endocrine disruptors; Plasticizers; Phthalates.

Graphical abstract



Pereyra-Camacho, M.A., Balderas-Hernández, V.E. & De Leon-Rodriguez, A. Biodegradation of diisononyl phthalate by a consortium of saline soil bacteria: optimisation and kinetic characterisation. *Appl Microbiol Biotechnol* (2021). <https://doi.org/10.1007/s00253-021-11255-5>

1 Introduction

Phthalic acid esters (PAEs) are a class of compounds employed as additives and plasticizers in several industrial applications and industrial products to improve their mechanical properties such as plasticity, strength and versatility (Xu et al. 2017; Li et al. 2019). Phthalate production has been reported to have reached 8 million tons worldwide (Jabesa and Ghosh 2016; Zhang et al. 2018). PAEs are referred as endocrine disruptors and their hepatotoxic, teratogenic and carcinogenic effects have been extensively documented (Katsikantami et al. 2016; Tang et al. 2016; Zhao et al. 2016). Thus, the U.S. Environmental Protection Agency and the European Union have listed phthalates as top priority contaminants (Zhang et al. 2018). From PAEs, the Di-isononyl phthalate (DINP) has become one of the most used plasticizers in the plastic industry. DINP is a mix of PAEs with 9-carbon alkyl chains of different lengths and branching distributions (Koch and Angerer 2007). DINP is used to replace di-(2-ethylhexyl) phthalate (DEHP) in several PVC products since it appears to have less toxic effects than DEHP (Hines et al. 2012; Chiang et al. 2020). However, other studies have found similar negative effects on reproduction when comparing DINP to DEHP (Chiang and Flaws 2019; Chiang et al. 2020). Therefore, DINP was classified as priority pollutant and endocrine-disruptor compound as well.

Since phthalates are not chemically bound to plastics, they can be released into different environmental matrices during their production and use (Jin et al. 2013; Zhao et al. 2016). Consequently, phthalates are found in several environmental matrices such as soil, water bodies, sediments, landfill leachates, and even in the

food and the atmosphere (Maitra 2016; Feng et al. 2018; Li et al. 2019). Concentrations of up to 500 mg L⁻¹ of PAEs have been reported in industrial wastewater (Ahmadi et al. 2017).

Phthalates can be degraded by abiotic processes like photolysis and chemical hydrolysis (Jin et al. 2013; Boll et al. 2020). However, microbial biodegradation is the best-known process for degrading PAEs from the environment. Several bacteria able to degrade phthalates isolated from different environmental matrices have been reported (Benjamin et al. 2015; Wang et al. 2017). They have the ability to use PAEs as only carbon source and energy, acting alone or as consortium (Ren et al. 2018). The most reported genera are: *Acinetobacter* sp., *Arthrobacter* sp., *Pseudomonas* sp., *Gordonia* sp., *Agrobacterium* sp., *Bacillus* sp., *Rhodococcus* sp., *Enterobacter* sp., *Burkholderia* sp., *Cyanobacteria*, among others (Babu and Wu 2010; Wu et al. 2010; Wang et al. 2012; Jin et al. 2013; Jin et al. 2015a; Liu et al. 2020). Existing evidence indicates that degradation and mineralization is faster using mixed cultures rather than monoaxenic cultures (He et al. 2013), because complete mineralization of PAEs involves a broad diversity of enzymes and cellular activities. In the environment, the complete degradation of phthalates is mainly carried out by the metabolic synergy of several microorganisms (Wu et al. 2010; Gao and Wen 2016).

In this study, a consortium of saline soil bacterial (SSB) was designed from bacterial isolates that were recognized as having the potential to use PAEs as only carbon source. The optimal culture conditions such as temperature, pH, and initial

DINP concentration for DINP degradation by the SSB-consortium were evaluated using the response surface methodology. Also, the kinetic characterization and degradation pathways of DINP were proposed from the intermediates detected by gas chromatography/mass spectrometry (GC/MS).

2 Hypothesis and aims

2.1 Hypothesis

Bacteria isolated from extreme environments have a robust metabolic machinery that will allow them to tolerate and degrade organic pollutants.

The use of bacterial consortia as a biological system for treatment of organic pollutants such as Diisononyl phthalate will promote an improvement in their degradation and transformation at laboratory level.

2.2 Aims

2.2.1 General

To isolate, identify and characterize diisononyl phthalate (DINP) degrading bacteria, as well as the optimization of the culture conditions for DINP degradation.

2.2.2 Specific

- a) To Identify degrading phthalates microorganisms from 2 bacterial collections: heavy metals tolerant and salinity-tolerant.
- b) To optimize the culture conditions for DINP biodegradation (temperature, pH and initial concentration of DINP).

- c) To identify the metabolic intermediates of DINP degradation and propose the bacterial biodegradation pathways.

3 Materials and Methods

3.1 Chemicals

DINP was acquired from Sigma-Aldrich (USA) with purity >99%. The stock of DINP (10 g L⁻¹) and Tween[®] 20 (10 g L⁻¹) were prepared in methanol and water, respectively, and kept at 4°C. Methanol and dichloromethane were HPLC grade (Sigma-Aldrich, Inc.). Chemicals used to prepare the culture media were analytical reagent grade and purchased from Sigma-Aldrich, Inc.

3.2 Bacterial collection and formulation of DINP-degrading SSB-consortium

21 bacterial isolates were obtained from soil samples from a Saltworks site (Zancarrón community, Santo Domingo municipality, San Luis Potosi, Mexico. 23° 23'40", 102° 11'46.5").

The identification of DINP tolerant isolates was evaluated using the spot-plate screening technique. DINP tolerance test was performed in solid mineral salt medium (MSM) plus DINP at 0, 500 or 1000 mg L⁻¹. The MSM contained per liter: CaCl₂ 0.02 g, (NH₄)₂SO₄ 2.0 g, MgSO₄ 0.16 g, KH₂PO₄ 4.5 g, K₂HPO₄ 5.85, 1 mL of trace elements solution (MgSO₄ 0.010 g, Na₂MoO₄•2H₂O 0.00036 g, Na₂SeO₃ 0.0002 g, NiCl₂•6H₂O 0.00034 g, FeCl₃•4H₂O 0.015 g, CoCl₂•6H₂O 0.0007 g, CuCl₂•2H₂O 0.0002 g, ZnCl₂ 0.023 g, MnCl₂•4H₂O 0.030 g) and agar 15 g. The medium was supplemented with yeast extract (0.3%) as a co-substrate to stimulate

the bacterial growth. Two microliters of each bacterial isolate were spot-cultivated onto MSM agar plate and incubated at 28°C for 72 hours. Bacteria able to grow at 1000 mg L⁻¹ of DINP were chosen as DINP-tolerant isolates and afterwards employed to formulate the SSB-consortium for further experiments.

3.3 Enrichment culture of the SSB-consortium

The enrichment culture technique for DINP degradation was similar to that described by Wu et al. (2010). Initially, the enrichment culture of the SSB-consortium was performed in a 50 mL Erlenmeyer flask with 25 mL of MSM plus 100 mg L⁻¹ DINP. Then, the culture was serially transferred to fresh medium supplemented with 100, 250, 500, 750 or 1000 mg L⁻¹ of DINP as only carbon substrate. Tween[®] 20 (500 mg L⁻¹) was added to improve the solubility of DINP in the medium (Jin et al. 2015a; Xu et al. 2017). All transfer cultures were initiated at an optical density at 595 nm (OD₅₉₅) of 0.1 and incubated for 5 days at 31°C and 180 rpm. The culture that grew and tolerated the highest concentration of DINP was selected as the final enrichment and used for the subsequent biodegradation experiments. The SSB-consortium is available upon request to the corresponding author.

3.4 Taxonomic identification of the SSB-consortium (16S Illumina sequencing)

The community analysis of the SSB-consortium was conducted through a 16S microbiota study. The total genomic DNA was extracted with Wizard[®] Genomic DNA Purification Kit. The V3 hypervariable region of the 16S rRNA gene was amplified by PCR using the V3-388F and V3-533R primers (Huse et al. 2008) and

Illumina adapters in the MiniSeq System (Illumina, Inc.) under standard conditions (300 cycles, 2x150 pair-end). The DNA extraction, amplification by PCR and data processing were carried out at the Microbial Genomics Laboratory (CIAD- Unit for Aquaculture, Mazatlán, Sinaloa, Mexico) (García-López et al. 2020). The sequencing data have been deposited in NCBI BioProject PRJNA691434 with the SRA submission accession SRR13415824 (**Table 1**).

Table 1. Bacterial diversity of the SSB-consortium sample for NCBI BioProject PRJNA691434

BioSample	Bacteria	Relative Abundance (%)	# Spots	% GC content	NCBI SRA accession
SSB-consortium ID: SAMN17293086	<i>Serratia</i> sp.	60.91%	14,553	52.68	SRR13415824
	<i>Methylobacillus</i> sp.	28.23%			
	<i>Achromobacter</i> sp.	7.05%			
	<i>Pseudomonas</i> sp.	3.11%			
	<i>Stenotrophomonas</i> sp.	0.45%			
	<i>Methyloversatilis</i> sp.	0.11%			
	<i>Delftia</i> sp.	0.07%			
	<i>Brevundimonas</i> sp.	0.07%			

3.5 Optimization of DINP biodegradation conditions

RSM analysis based on a Central Composite Design (CCD), was used to optimise the temperature, pH and initial concentration of DINP to improve the percentage of degradation of DINP by the SSB-consortium. The values of the independent variables are shown in **Table 2**.

Table 2. Levels of independent variables used in the CCD for the optimization of DINP degradation.

Independent variables (units)	Symbol	Code levels of variables				
		- α	-1	0	1	+ α
Temperature (°C)	X ₁	21	25	31	37	41
pH	X ₂	5.3	6	7	8	8.6
DINP concentration (mg L ⁻¹)	X ₃	79.5	250	500	750	920.4

A quadratic polynomial model (**Eq. 1**) was suggested to explain the effects of independent variables on the dependent variable based on the data obtained in the experimental design.

$$Y_i = \beta_0 + \sum \beta_i X_i + \sum \beta_{ii} X_i^2 + \sum \beta_{ij} X_i X_j \quad (1)$$

Where Y_i is the corresponding response, β_0 is the intercept term, β_i is the linear coefficient, β_{ii} is the squared coefficient and β_{ij} is the interaction coefficient, whereas X_i and X_j are the variables (Zhao et al. 2018a).

The experimental design and analysis of variance (ANOVA) was employed to determine the interaction between independent variables (temperature, pH and DINP concentration) and the dependent variable. The statistical analysis and response surface analysis were performed with Design-Expert[®] Software (Minneapolis, MN).

3.6 Biodegradation kinetics of DINP by the SSB-consortium

The SSB-consortium was inoculated in MSM plus 500 mg L⁻¹ DINP at 31°C and 180 rpm for 72 h. Subsequently, the bacteria were collected and washed two times with PBS 0.1 mM (pH 7.0) and resuspended in MSM. Cell suspension with an OD₅₉₅ of 0.1 was employed to inoculate the medium for degradation assays.

Biodegradation kinetics were carried out in 200 mL MSM plus 500 mg L⁻¹ DINP as the only carbon source under the optimum culture conditions for 168 h. Tween[®] 20 (500 mg L⁻¹) was added to enhance the solubility of DINP (Jin et al. 2015a; González et al. 2020). Non-inoculated culture was used as control. Samples were collected every 24 h to quantify the residual DINP and cell growth was followed as optical density at OD₅₉₅. Residual DINP in the samples was extracted as previously reported by Zhang et al.,(2018) and measured by GC/MS. Experiments were carried out in triplicates.

The biodegradation of DINP (%) was determined from the difference in concentration in the medium according to **Eq. 2**:

$$\text{DINP Biodegradation (\%)} = (C_0 - C_t)/C_0 * 100 \quad (2)$$

Where C₀ is the initial concentration of DINP in the medium, C_t is the final concentration of DINP at time interval t (Zhao et al. 2018a).

To evaluate the influence of the initial concentration of DINP on its degradation kinetics, different concentrations (250, 500, 750 mg L⁻¹) were evaluated and fixed to a first-order kinetic model **Eq. 3**:

$$\ln C = -kt + A \quad (3)$$

Where C is the concentration of DINP (mg L⁻¹), k is the first-order constant (h⁻¹), t is the time (h), and A is a constant (Zhang et al. 2018; Zhao et al. 2018a).

Then the half-life time (t_{1/2}) of the DINP biodegradation was evaluated according to the following **Eq. 4**:

$$t_{1/2} = \ln 2/k \quad (4)$$

3.7 Analytical methods

Optical density at 595 nm (OD₅₉₅) was measured with a microplate reader (iMark™, Bio-Rad Laboratories, Inc.). For residual DINP quantification and identification of its metabolites, 10 mL of culture were collected from the culture media daily during 168 h of incubation. Then, compounds were extracted with dichloromethane from the samples as reported by Zhang et al., (2018). The organic phase was collected and evaporated to dryness and resuspended in 1 mL of methanol. DINP quantification and metabolite identification were performed by GC/MS 7820A/5977E (Agilent Technologies, Inc.) using a HP-5MS column (30 m x 250 μm x 0.25 μm) with helium as carrier gas at a flow rate of 1.0 mL min⁻¹. The chromatographic conditions reported by González et al. 2020 were used. The

identification of the metabolites was done by comparing their mass spectra to those obtained in the NIST 14 library of the MS database (Tang et al. 2016).

4 Results

4.1 Identification and characterization of the SSB-consortium.

Twenty-one bacteria from a collection of saltwork site-isolates were cultured in MSM-agar plates at high concentrations of DINP (500-1000 mg L⁻¹) using yeast extract as co-substrate. From those, 18 bacteria grew in 500 mg L⁻¹ DINP, and only 11 grew with 1000 mg L⁻¹ DINP. For the design of the SSB-consortium, only the microorganisms that grew on and tolerated 1000 mg L⁻¹ DINP were selected. When DINP was the only source of carbon in the culture medium of the enrichment assay, the bacterial growth capacity and tolerance decreased. Previous reports have indicated that the use of co-substrates, such as yeast extract, in the culture medium could promote and enhance the cell growth and phthalate degradation (Li et al. 2019). After several transfers of the enrichment culture using fresh medium, a consortium capable of growing up to 500 mg L⁻¹ of DINP was obtained and used for further DINP biodegradation experiments.

Bacteria present in the SSB-consortium were taxonomically identified through microbiome 16S rRNA sequencing (**Fig. 1**). The results indicated that seven families compose the bacterial community of the SSB-consortium, the main families were *Enterobacteriaceae* (60.91%), *Methylophilaceae* (28.23%), *Burkholderiaceae* (7.12%), *Pseudomonadaceae* (3.11%), representing 99.37% of

the reads. The families *Xanthomonadaceae* (0.45%), *Rhodocyclaceae* (0.11%), and *Caulobacteraceae* (0.07%), were also found but at relatively low abundance. The analysis at genus level revealed that the SSB-consortium is composed by 8 bacterial genera, based on their relative abundance the most predominant genera were *Serratia* sp. (60.91%), *Methylobacillus* sp. (28.23%), *Achromobacter* sp. (7.05%) and *Pseudomonas* sp. (3.11%). The relative abundance of *Stenotrophomonas* sp., *Methyloversatilis* sp., *Delftia* sp. and *Brevundimonas* sp. were 0.45%, 0.11%, 0.07%, 0.07%, respectively.




Figure 1. Identification of the bacterial diversity of the SSB-consortium. (a) at family level; (b) at genus level.

4.2 Optimization of operational conditions for DIMP biodegradation

Based on a Central Composite Design a RSM analysis was used to assess and optimize the influence of the cultivation variables: temperature (X_1), pH (X_2) and initial DIMP concentration (X_3) on the DIMP biodegradation. The results of the % of DIMP degradation obtained by the CCD are shown in **Table 3**. From the regression

analysis of the experimental data a quadratic polynomial model was obtained to describe the degradation of DINP (Eq. 5).

$$Y_{DINP} = 97.44 - 11.71X_1 + 2.33X_2 - 23.48X_3 - 4.26X_1X_2 + 19.13X_1X_3 - 4.26X_2X_3 - 31.94X_1^2 - 32.94X_2^2 - 15.82X_3^2 \quad (5)$$

Where Y is the predicted % biodegradation of DINP by the SSB-consortium, X_1 , X_2 , X_3 are the coded variables that correspond to temperature, pH and initial DINP concentration, respectively.

Table 4 shows the ANOVA analysis performed to assess the importance of the adjusted model together with the effects of the variables with their linear, quadratic, and interactive terms associated with the degradation of DINP. Moreover, to verify the suitability of the model and the statistical significance of the terms, a p -value <0.05 , was used. The model p -value of 0.0006 indicated that the model was statistically significant (Cisneros de la Cueva et al. 2017). The accuracy and reliability of the model were assessed according to the determination coefficient ($R^2= 0.9567$), indicating that the DINP biodegradation model could explain the 95.67% variability of the dependent variable, and the good fit between the model and the experimental biodegradation data, indicating that the treatment was highly significant.

Table 3. Matrix of the Composite Central Design and the response: % degradation of DINP.

Run	Levels of independent variables			Response
	Temperature (°C)	pH	Concentration of DINP (mg L ⁻¹)	DINP degradation (%)
1	25	6	250	62.30
2	37	8	250	2.00
3	25	6	750	0.67
4	31	7	500	97.60
5	37	6	250	0.80
6	25	8	250	95.28
7	31	7	500	97.30
8	37	6	750	0.93
9	25	8	750	0.67
10	37	8	750	0.13
6	31	7	79.55	97.11
12	31	7	920.45	0.92
13	41.1	7	500	2.40
14	31	8.68	500	0.40
15	20.9	7	500	6.20
16	31	7	500	97.88
17	31	5.32	500	0.80
18	31	7	500	97.30

The percentage of biodegradation in the response surface graphs and contour graphs (**Fig. 2**) directly showed the influence of the initial concentration of DINP, temperature and pH. The results showed a degradation range from 0.13 to 97.88% (**Table 3**). The regression analysis indicated that linear and quadratic terms of temperature (X_1) and initial DINP concentration (X_3) had significant impact ($p < 0.05$) in the degradation of DINP, while the linear term of pH (X_2) and their interaction terms (X_1X_2 and X_2X_3) had no significant effects ($p > 0.05$) as shown in

Table 4, indicating that the pH (in the evaluated range) was not a key factor in the biodegradation of DINP. The maximum theoretical DINP degradation was 97.52% and the optimal culture conditions were pH 7.0, temperature of 31°C, and the initial concentration of 500 mg L⁻¹ DINP (**Table 3**).

A final set of DINP biodegradation experiments using the predicted optimal culture conditions was implemented. DINP degradation was 98.88%, which was similar to that predicted by the model, demonstrating the suitability of the obtained model to optimize the conditions of DINP biodegradation. Maximum % of DINP biodegradation was reached after 168 h of cultivation.

The effect of the variables pH, temperature and initial DINP concentration on the % of degradation was evaluated using ANOVA (**Table 4**). Temperature had a significant effect ($p < 0.0232$) and it was a key factor in the DINP degradation. The surface and contour plots (**Fig. 2 a-d**) show the interaction of temperature with pH and the initial concentration of DINP, indicating that temperature had an important role in the biodegradation of DINP. At 25°C, 20-30% of DINP degradation were observed, whereas at 37°C there was no degradation. The pH in the range of 6.0-8.0 was not statistically significant ($p > 0.5829$), indicating that it did not play an important role in DINP biodegradation, thus a pH 7.0 was chosen as the optimal condition.




Figure 2. Contour and 3D response surface plots showing the influence of pH, temperature, and initial DINP concentration on the % of DINP biodegradation by the SSB-consortium.

On the other hand, the initial concentration of DINP showed to be statistically significant ($p < 0.0007$), the results showed that the SSB-consortium had the capacity to degrade up to 500 mg L⁻¹ of DINP, whereas higher concentrations inhibited the growth of the consortium.

Table 4. ANOVA of the % of DINP degradation obtained by Central Composite Design.

Source	Sum of Squares	Df	Mean Square	F Value	p-value
Block	18.10	1	18.10		
Model	34549.43	9	3838.83	17.17	0.0006
X ₁ -Temperature	1873.36	1	1873.36	8.38	0.0232
X ₂ -pH	74.08	1	74.08	0.33	0.5829
X ₃ -DINP Concentration	7528.64	1	7528.64	33.68	0.0007
X ₁ X ₂	145.18	1	145.18	0.65	0.4468
X ₁ X ₃	2926.13	1	2926.13	13.09	0.0085
X ₂ X ₃	145.18	1	145.18	0.65	0.4468
X ₁ ²	12574.78	1	12574.78	56.25	0.0001
X ₂ ²	13636.53	1	13636.53	61.00	0.0001
X ₃ ²	3145.43	1	3145.43	14.07	0.0072
Residual	1564.82	7	223.55		
Lack of Fit	1564.61	5	312.92	2935.48	0.0003
Pure Error	0.210	2	0.110		
Cor Total	36132.35	17			

4.3 Biodegradation kinetics of DINP

To determine the kinetic parameters of DINP biodegradation by the SSB-consortium, a set of cultures were carried out in MSM under optimal culture conditions of 31°C, pH 7, and 500 mg L⁻¹ DINP (**Fig. 3**). Optical density increased from 0.1 to 1.73 ±0.06 in the presence of DINP in 144 h, while no increase was observed in the control cultures without the DINP (**Fig. 3a**). As observed in **Fig. 3b**, 99% of the DINP was degraded in 144 h in the inoculated cultures, whereas no change of DINP concentration was observed in the cell-free cultures. These results indicated that the SSB-consortium used DINP as the only source of carbon and energy, to confirm this asseveration, additional experiments by inoculating the SSB-consortium in MSM without DINP, but containing Tween 20 were carried out, and no growth was observed after 168 h of cultivation. Tween 20 did not show toxicity at the used concentration (500 mg L⁻¹), on the other hand it helped in the assimilation of the DINP by its emulsifying property, as previously reported (Jin et al. 2015a; Xu et al. 2017).

To evaluate the biodegradation rate of DINP by the SSB-consortium, DINP was used at different initial concentrations (250, 500 and 750 mg L⁻¹) in MSM. As shown in **Fig. 4**, DINP biodegradation followed a first-order kinetics and the parameters are summarized in **Table 5**. In the cultures with the initial concentration of 750 mg L⁻¹ DINP, no degradation was observed, indicating an inhibitory effect. DINP was rapidly biodegraded when the initial concentration was 250 mg L⁻¹ compared to 500 mg L⁻¹ (**Fig. 4**). First-order degradation constants (*k*) were 0.0629 h⁻¹ and 0.0543 h⁻¹, whereas the half-life times (*t*_{1/2}) were 11.01 and 12.76 h for 250

and 500 mg L⁻¹, respectively. The $t_{1/2}$ corresponds to the time it takes to decrease the DINP concentration in half (Benjamin et al. 2015; Zhao et al. 2018a).




Figure 3. Growth of the SSB-consortium and DINP biodegradation kinetics started at 500 mg L⁻¹ DINP: a) Evaluation of cell growth in MSM with or without DINP as the only carbon source, and b) degradation of DINP by the SSB-consortium under optimal conditions (31 °C, pH 7 and 500 mg L⁻¹) in MSM. Non-inoculated MSM was employed as control (black squares). Error bars indicate the standard error of the three replicates.




Figure 4. Biodegradation of DINP at different initial concentrations performed by the SSB-consortium in MSM at 31°C and pH 7. DINP concentrations used were: ▲; 250 mg L⁻¹, ●; 500 mg L⁻¹ and ○; 750 mg L⁻¹.

Table 5. Kinetic equations of DINP biodegradation at different initial DINP concentrations.

Initial Concentration (mg L ⁻¹)	Kinetic equations	t _{1/2} (h)	R ²
250	Ln C = -0.0629t + 7.6241	11.01	0.997
500	Ln C = -0.0543t + 10.9140	12.76	0.987
750	-	-	-

4.4 Identification of DINP degradation intermediates and metabolic pathways

To determine the biodegradation pathways of DINP by the SSB-consortium, their intermediates were identified by GC/MS. A total of four intermediates were detected (**Fig. 5** and **Table 6**). At the beginning of the cultivation only DINP (retention time, RT: 9.6 – 10.6 min) was detected (**Fig. 5a**). Throughout the time of cultivation, DINP decreased and at the same time, 4 new compounds were detected at the 72 h of culture (**Fig. 5b**). These compounds were monoisononyl phthalate (MiNP; RT: 8.0 - 8.5 min), methyl nonyl phthalate (MNP; RT: 8.11 min), iso-nonanol (INA; RT: 3.8 - 4.5 min), and dimethyl phthalate (DMP; RT: 5.68 min). Where MiNP, MNP and INA were the most abundant. As observed in **Fig. 5c** the relative abundance of metabolites continued gradually changing and DINP was almost completely degraded at 120 h. MiNP and MNP decreased and the peaks of DMP and INA increased, showing the SSB-consortium was able to transform DINP

and its metabolites into more simple compounds. At the end of the culture (168 h) no peaks of DINP, MiNP, MNP were detected, indicating that they were completely degraded, and only traces of Iso-nonanol and DMP remained (**Fig. 5d**). No intermediates were detected in the control samples (without the inoculum) and only DINP was detected.

Based on the metabolites detected by GC/MS, β -oxidation also occurred as a simultaneous pathway in the first stage of the biodegradation. DINP was transformed into MiNP and INA by de-esterification, whereas MNP and DMP were produced by β -oxidation. Then, the products were degraded into phthalic acid (PA). Since PA was not observed, it could indicate that it was quickly incorporated and metabolized into the central carbon pathways to CO₂ and H₂O (**Fig. 6**). Li et al., 2018 reported a similar effect during the biodegradation of DEHP with the consortium LF, phthalic acid was not detected.

Table 6. Identification of DINP biodegradation metabolites by GC/MS*.

Metabolites	Molecular ion (<i>m/z</i>)	Retention Time
		RT (min)
Diisononyl phthalate	149, 280.9, 135	9.6 -10.6
Monoisononyl phthalate	149, 163, 150	8.0 - 8.5
Methyl nonyl phthalate	163, 149, 181	8.118
Iso-nonanol	56, 43, 70	3.8 - 4.5
Dimethyl phthalate	163, 77, 92	5.692

*GC Column HP-5MS (30 m x 250 μ m x 0.25 μ m).




Figure 5. GC/MS chromatograms of DINP biodegradation by the SSB-consortium:

a) 0 h, b) 72 h, c) 120 h, and d) 168 h of bacterial growth.




Figure 6. Proposed pathways for DINP degradation by the SSB-consortium. The dotted box indicates that the metabolite was not detected in this study.

5 Discussion

In this study, an SSB-consortium formulated with bacterial isolates from a saltworks site, was able to degrade DINP and use it as the only carbon and energy source. SSB-consortium was mainly constituted by four bacterial genera *Serratia* sp., *Methylobacillus* sp., *Achromobacter* sp. and *Pseudomonas* sp. Nowadays, there are many reports on the ability of *Achromobacter* sp. and *Pseudomonas* sp. to degrade organic contaminants, including phthalate esters, bisphenol A, phenols, and other aromatic compounds (Xu et al. 2005; Benjamin et al. 2015; Jin et al. 2015a; Liu et al. 2015; Li et al. 2018). However, there are few reports on degradation of phthalates by the genera *Serratia* sp. (Chuang Li 2012) and *Methylobacillus* sp. (Maitra 2016), which represent the higher percentage of relative abundance in the consortium. Previous studies indicate that the use of bacterial consortia is more suitable for bioremediation than pure bacterial strains

(Wang et al. 2017; Li et al. 2018), indicating that the SSB-consortium is an ideal candidate for bioremediation of sites contaminated by phthalate esters.

Results showed that temperature and initial DINP concentration play a significant role on the degradation process, while pH (range evaluated) was not a key factor in the biodegradation of DINP. Several studies noted a similar statistical effect, indicating that bacteria may be able to adapt and degrade over wide pH ranges (Xu et al. 2005; Li et al. 2019). The biodegradation results obtained by the SSB-consortium are promising, so far, there is only one report of DINP biodegradation by Park et al., (2008). They reported 100% degradation of 500 mg L⁻¹ DINP in 10 days by *Sphingobium chungbukense*, while the consortium used here, attained total degradation in less than 7 days. Optimal conditions reported were similar in both studies (pH 7.0), and temperature of 30 and 31°C, respectively. These results suggest that the use of bacterial consortia is more efficient for the DINP degradation.

The results for DINP biodegradation kinetic indicated the SSB-consortium used DINP as the sole source of carbon and energy and followed a first-order kinetics. The ability of the SSB-consortium was outstanding, to biodegrade DINP (500 mg L⁻¹), resulting in an almost complete elimination (99%) in 168 h with a k of 0.0543 h⁻¹, whereas the half-life time ($t_{1/2}$) was 12.76 h. These results are comparable with previous studies reporting the degradation of phthalates with long and short alkyl-chains. For instance, Li et al., (2018) found that the LF consortium could degrade 500 mg L⁻¹ DEHP completely with a k of 0.0404 h⁻¹, while the SSB-consortium

degraded the same amount of DINP with a higher degradation constant (0.0543 h^{-1}), suggesting that the SSB-consortium is more efficient than consortium LF. Zhang et al., (2018) described the completely degradation of 500 mg L^{-1} DEHP by *Bacillus mojavensis* B1811 with a k of 0.0149 h^{-1} . Lu et al., (2009) reported a degradation constant of 0.0062 h^{-1} and $t_{1/2}$ of 4.63 days for a mixture of PAEs (di-n-butyl phthalate, diethyl phthalate and dimethyl phthalate) at a concentration of 600 mg L^{-1} using a *Rhodococcus* sp. L4. Feng et al., (2002) described the degradation of DEHP by *Pseudomonas fluorescens* FS1, where they found that the k was 0.0408 h^{-1} . These results showed that the SSB-consortium could biodegrade DINP more efficiently compared to the other systems, being a viable alternative for bioremediation of phthalates with long alkyl-chains such as DINP.

The presence of MiNP and iso-nonanol suggests that de-esterification was carried out, whereas the presence of MNP indicates that the biodegradation of DINP also occurred through β -oxidation. One of the main steps in the degradation of phthalates involves the hydrolysis of the ester bonds through esterases, resulting in generation of the monoesters that are subsequently hydrolysed to phthalic acid (PA) and alcohols (Liang et al. 2008; Zhang et al. 2018; Li et al. 2019). Both pathways (β -oxidation and de-esterification) in the biodegradation of long alkyl-chains phthalates have been described previously (Benjamin et al. 2015; Ren et al. 2018; Li et al. 2019; Bai et al. 2020). To our knowledge, only one de-esterification had been proposed for the biodegradation of DINP by *Sphingobium chungbukense* (Park et al. 2008), whereas in this work, DINP degradation by the SSB-consortium was performed by de-esterification and β -oxidation simultaneously. This may be

due to the use of bacterial consortia, previous studies indicated that the use of consortiums improve the degradation efficiency of environmental pollutants (He et al. 2013; Wang et al. 2017; Li et al. 2018). Thus, SSB-consortium degraded the DINP faster than that reported using a single bacterium.

6 Conclusion

In conclusion, the results obtained in this research demonstrate that the SSB-consortium degrades up to 500 mg L⁻¹ of DINP following a first-order kinetics and the temperature and the initial concentration of DINP are the main factors affecting its biodegradation without accumulating toxic by-products. Therefore, the SSB-consortium is a potential candidate to bioremediate effluents contaminated by DINP.

7 References

- Abraham J, Silambarasan S (2015) Plant Growth Promoting Bacteria *Enterobacter asburiae* JAS5 and *Enterobacter cloacae* JAS7 in Mineralization of Endosulfan. *Appl Biochem Biotechnol* 175:3336–3348 . <https://doi.org/10.1007/s12010-015-1504-7>
- Ahmadi E, Yousefzadeh S, Ansari M, Ghaffari HR (2017) Performance , kinetic , and biodegradation pathway evaluation of anaerobic fixed film fixed bed reactor in removing phthalic acid esters from wastewater. *Nat Publ Gr* 1–14 . <https://doi.org/10.1038/srep41020>
- Al-Saleh I, Elkhatib R, Al-Rajoudi T, Al-Qudaihi G (2017) Assessing the concentration of phthalate esters (PAEs) and bisphenol A (BPA) and the genotoxic potential of treated wastewater (final effluent) in Saudi Arabia. *Sci Total Environ* 578: . <https://doi.org/10.1016/j.scitotenv.2016.10.207>

- Babu B, Wu JT (2010) Biodegradation of phthalate esters by cyanobacteria. *J Phycol* 46:1106–1113 . <https://doi.org/10.1111/j.1529-8817.2010.00896.x>
- Bai N, Li S, Zhang J, Zhang H, Zhang H, Zheng X, Lv W (2020) Efficient biodegradation of DEHP by CM9 consortium and shifts in the bacterial community structure during bioremediation of contaminated soil. *Environ Pollut* 266:115112 . <https://doi.org/10.1016/j.envpol.2020.115112>
- Benjamin S, Masai E, Kamimura N, Takahashi K, Anderson RC, Faisal PA (2017) Phthalates impact human health: Epidemiological evidences and plausible mechanism of action. *J Hazard Mater* 340:360–383 . <https://doi.org/10.1016/j.jhazmat.2017.06.036>
- Benjamin S, Pradeep S, Sarath Josh M, Kumar S, Masai E (2015) A monograph on the remediation of hazardous phthalates. *J Hazard Mater* 298:58–72 . <https://doi.org/10.1016/j.jhazmat.2015.05.004>
- Boll M, Geiger R, Junghare M, Schink B (2020) Microbial degradation of phthalates: biochemistry and environmental implications. *Environ Microbiol Rep* 12:3–15 . <https://doi.org/10.1111/1758-2229.12787>
- Chiang C, Flaws JA (2019) Subchronic Exposure to Di(2-ethylhexyl) Phthalate and Diisononyl Phthalate during Adulthood Has Immediate and Long-Term Reproductive Consequences in Female Mice. *Toxicol Sci*. <https://doi.org/10.1093/toxsci/kfz013>
- Chiang C, Lewis LR, Borkowski G, Flaws JA (2020) Late-life consequences of short-term exposure to di(2-ethylhexyl) phthalate and diisononyl phthalate during adulthood in female mice. *Reprod Toxicol*. <https://doi.org/10.1016/j.reprotox.2019.12.006>
- Chuang Li (2012) Biodegradation of an endocrine-disrupting chemical di-n-butyl phthalate by *Serratia marcescens* C9 isolated from activated sludge. *African J Microbiol Res*. <https://doi.org/10.5897/ajmr11.1177>
- Cisneros de la Cueva S, Alvarez Guzmán CL, Balderas Hernández VE, De León Rodríguez A (2017) Optimization of biohydrogen production by the novel psychrophilic strain N92 collected from the Antarctica. *Int J Hydrogen Energy* 43:13798–13809 . <https://doi.org/10.1016/j.ijhydene.2017.11.164>

- Dangi AK, Sharma B, Hill RT, Shukla P (2019) Bioremediation through microbes: systems biology and metabolic engineering approach. *Crit Rev Biotechnol* 39:79–98 . <https://doi.org/10.1080/07388551.2018.1500997>
- Das MT, Kumar SS, Ghosh P, Shah G, Malyan SK, Bajar S, Thakur IS, Singh L (2021) Remediation strategies for mitigation of phthalate pollution: Challenges and future perspectives. *J. Hazard. Mater.* 409:124496
- Feng NX, Feng YX, Liang QF, Chen X, Xiang L, Zhao HM, Liu BL, Cao G, Li YW, Li H, Cai QY, Mo CH, Wong MH (2021a) Complete biodegradation of di-n-butyl phthalate (DBP) by a novel *Pseudomonas* sp. YJB6. *Sci Total Environ* 761: . <https://doi.org/10.1016/j.scitotenv.2020.143208>
- Feng NX, Feng YX, Liang QF, Chen X, Xiang L, Zhao HM, Liu BL, Cao G, Li YW, Li H, Cai QY, Mo CH, Wong MH (2021b) Complete biodegradation of di-n-butyl phthalate (DBP) by a novel *Pseudomonas* sp. YJB6. *Sci Total Environ* 761:143208 . <https://doi.org/10.1016/j.scitotenv.2020.143208>
- Feng NX, Yu J, Mo CH, Zhao HM, Li YW, Wu BX, Cai QY, Li H, Zhou DM, Wong MH (2018) Biodegradation of di-n-butyl phthalate (DBP) by a novel endophytic *Bacillus megaterium* strain YJB3. *Sci Total Environ* 616–617:117–127 . <https://doi.org/10.1016/j.scitotenv.2017.10.298>
- Feng Z, Kunyan C, Jiamo F, Gouying S, Huifang Y (2002) Biodegradability of di(2-ethylhexyl) phthalate by *Pseudomonas fluorescens* FS1. *Water Air Soil Pollut* 140:297–305
- Gao D, Li Z, Wang H, Liang H (2018) An overview of phthalate acid ester pollution in China over the last decade: Environmental occurrence and human exposure. *Sci. Total Environ.* 645:1400–1409
- Gao DW, Wen ZD (2016) Phthalate esters in the environment: A critical review of their occurrence, biodegradation, and removal during wastewater treatment processes. *Sci. Total Environ.*
- García-López R, Cornejo-Granados F, Lopez-Zavala AA, Sánchez-López F, Cota-Huizar A, Sotelo-Mundo RR, Guerrero A, Mendoza-Vargas A, Gómez-Gil B, Ochoa-Leyva A (2020) Doing more with less: A comparison of 16S hypervariable regions in search of defining the shrimp microbiota.

- Microorganisms. <https://doi.org/10.3390/microorganisms8010134>
- Giuliani A, Zuccarini M, Cichelli A, Khan H, Reale M (2020) Critical review on the presence of phthalates in food and evidence of their biological impact. *Int J Environ Res Public Health* 17:1–43 . <https://doi.org/10.3390/ijerph17165655>
- González JL, Marco E, Camacho AP, León A De, Alicia R, Lagunes G (2020) Biodegradation of recalcitrant compounds and phthalates by culturable bacteria isolated from *Liometopum apiculatum* microbiota. *World J Microbiol Biotechnol* 1: . <https://doi.org/10.1007/s11274-020-02850-1>
- He Z, Xiao H, Tang L, Min H, Lu Z (2013) Bioresource Technology Biodegradation of di-n-butyl phthalate by a stable bacterial consortium , HD-1 , enriched from activated sludge. *128:526–532*
- Hines CJ, Hopf NB, Deddens JA, Silva MJ, Calafat AM (2012) Occupational exposure to diisononyl phthalate (DiNP) in polyvinyl chloride processing operations. *Int Arch Occup Environ Health* 85:317–325 . <https://doi.org/10.1007/s00420-011-0674-z>
- Huang H, Zhang XY, Chen TL, Zhao YL, Xu DS, Bai YP (2019) Biodegradation of Structurally Diverse Phthalate Esters by a Newly Identified Esterase with Catalytic Activity toward Di(2-ethylhexyl) Phthalate. *J Agric Food Chem* 67:8548–8558 . <https://doi.org/10.1021/acs.jafc.9b02655>
- Huang X-J, Du H, Deng X-L, Chen Y-H, Xiang L, Li Y-W, Li H, Mo C-H, Cai Q-Y, Zhao H-M (2021) New insights into the evolution of bacterial community during the domestication of phthalate-degrading consortium. *J Clean Prod* 303:127064 . <https://doi.org/10.1016/j.jclepro.2021.127064>
- Huse SM, Dethlefsen L, Huber JA, Welch DM, Relman DA, Sogin ML (2008) Exploring microbial diversity and taxonomy using SSU rRNA hypervariable tag sequencing. *PLoS Genet*. <https://doi.org/10.1371/journal.pgen.1000255>
- Jabesa A, Ghosh P (2016) Removal of diethyl phthalate from water by ozone microbubbles in a pilot plant. *J Environ Manage*. <https://doi.org/10.1016/j.jenvman.2016.05.072>
- Jin D, Bai Z, Chang D, Hoefel D, Jin B, Wang P, Wei D, Zhuang G (2012) Biodegradation of di-n-butyl phthalate by an isolated *Gordonia* sp. strain QH-

- 11: Genetic identification and degradation kinetics. *J Hazard Mater* 221–222:80–85 . <https://doi.org/10.1016/j.jhazmat.2012.04.010>
- Jin D, Kong X, Cui B, Bai Z, Zhang H (2013) Biodegradation of di-n-butyl phthalate by a newly isolated halotolerant *Sphingobium* sp. *Int J Mol Sci* 14:24046–24054 . <https://doi.org/10.3390/ijms141224046>
- Jin D, Kong X, Li Y, Bai Z, Zhuang G, Zhuang X, Deng Y (2015a) Biodegradation of di-n-butyl phthalate by *Achromobacter* sp. Isolated from rural domestic wastewater. *Int J Environ Res Public Health* 12:13510–13522 . <https://doi.org/10.3390/ijerph121013510>
- Jin D, Kong X, Li Y, Bai Z, Zhuang G, Zhuang X, Deng Y (2015b) Biodegradation of di-n-butyl phthalate by *Achromobacter* sp. Isolated from rural domestic wastewater. *Int J Environ Res Public Health* 12:13510–13522 . <https://doi.org/10.3390/ijerph121013510>
- Katsikantami I, Sifakis S, Tzatzarakis MN, Vakonaki E, Kalantzi OI, Tsatsakis AM, Rizos AK (2016) A global assessment of phthalates burden and related links to health effects. *Environ Int* 97:212–236 . <https://doi.org/10.1016/j.envint.2016.09.013>
- Khadka S, Nshimiyimana JB, Zou P, Koirala N, Xiong L (2020) Biodegradation Kinetics of Diethyl Phthalate by Three Newly Isolated Strains of *Pseudomonas*. *Sci African* 8: . <https://doi.org/10.1016/j.sciaf.2020.e00380>
- Kim H, Nam KS, Oh S, Son S, Jeon D, Chan Gye M, Shin I (2019) Toxicological assessment of phthalates and their alternatives using human keratinocytes. *Environ Res* 175: . <https://doi.org/10.1016/j.envres.2019.05.007>
- Koch HM, Angerer J (2007) Di-iso-nonylphthalate (DINP) metabolites in human urine after a single oral dose of deuterium-labelled DINP. *Int J Hyg Environ Health*. <https://doi.org/10.1016/j.ijheh.2006.11.008>
- Kolb SA, O’Loughlin EJ, Gsell TC (2019) Characterization of phthalate-degrading bacteria from Asian carp microbiomes and riverine sediments. *Int Biodeterior Biodegrad*. <https://doi.org/10.1016/j.ibiod.2019.104727>
- Li F, Liu Y, Wang D, Zhang C, Yang Z, Lu S, Wang Y (2018) Biodegradation of di-(2-ethylhexyl) phthalate by a halotolerant consortium LF. *PLoS One*.

- <https://doi.org/10.1371/journal.pone.0204324>
- Li J, Zhang J, Yadav MP, Li X (2019) Biodegradability and biodegradation pathway of di-(2-ethylhexyl) phthalate by *Burkholderia pyrrocinia* B1213. *Chemosphere*. <https://doi.org/10.1016/j.chemosphere.2019.02.194>
- Liang DW, Zhang T, Fang HHP, He J (2008) Phthalates biodegradation in the environment. *Appl Microbiol Biotechnol* 80:183–198 . <https://doi.org/10.1007/s00253-008-1548-5>
- Liu F, Liu Q, Zhang Y, Liu Y, Wan Y, Gao K, Huang Y, Xia W, Wang H, Shi Y, Huang Z, Lu B (2015) Molecularly imprinted nanofiber membranes enhanced biodegradation of trace bisphenol A by *Pseudomonas aeruginosa*. *Chem Eng J* 262:989–998 . <https://doi.org/10.1016/j.cej.2014.10.046>
- Liu G, Zhang S, Yang K, Zhu L, Lin D (2016) Toxicity of perfluorooctane sulfonate and perfluorooctanoic acid to *Escherichia coli*: Membrane disruption, oxidative stress, and DNA damage induced cell inactivation and/or death. *Environ Pollut* 214:806–815 . <https://doi.org/10.1016/j.envpol.2016.04.089>
- Liu T, Li J, Qiu L, Zhang F, Linhardt RJ, Zhong W (2020) Combined genomic and transcriptomic analysis of the dibutyl phthalate metabolic pathway in *Arthrobacter* sp. ZJUTW. *Biotechnol Bioeng*. <https://doi.org/10.1002/bit.27524>
- Lu M, Jiang W, Gao Q, Zhang M, Hong Q (2020) Degradation of dibutyl phthalate (DBP) by a bacterial consortium and characterization of two novel esterases capable of hydrolyzing PAEs sequentially. *Ecotoxicol Environ Saf* 195: . <https://doi.org/10.1016/j.ecoenv.2020.110517>
- Lu Y, Tang F, Wang Y, Zhao J, Zeng X, Luo Q, Wang L (2009) Biodegradation of dimethyl phthalate, diethyl phthalate and di-n-butyl phthalate by *Rhodococcus* sp. L4 isolated from activated sludge. *J Hazard Mater* 168:938–943 . <https://doi.org/10.1016/j.jhazmat.2009.02.126>
- Maitra VKSS (2016) Biodegradation of endocrine disruptor dibutyl phthalate (DBP) by a newly isolated *Methylobacillus* sp . V29b and the DBP degradation pathway. *3 Biotech* 6:1–12 . <https://doi.org/10.1007/s13205-016-0524-5>
- Park JM, Jeon M, Lim ES, Um HJ, Kim YC, Min J, Kim YH (2008) Biodegradation

- of a phthalate plasticizer, di-isononyl phthalate (DINP), by *Sphingobium chungbukense*. *Curr Microbiol*. <https://doi.org/10.1007/s00284-008-9232-7>
- Patrón-Soberano A, Núñez-Luna BP, Casas-Flores S, De Las Peñas A, Domínguez-Espíndola RB, Rodríguez-González V (2017) Photo-assisted inactivation of: *Escherichia coli* bacteria by silver functionalized titanate nanotubes, Ag/H₂Ti₂O₅·H₂O. *Photochem Photobiol Sci* 16:854–860 . <https://doi.org/10.1039/c6pp00237d>
- Pereyra-camacho MA, Balderas-hernández VE, Leon-rodriguez A De (2021) Biodegradation of diisononyl phthalate by a consortium of saline soil bacteria : optimisation and kinetic characterisation
- Ren L, Lin Z, Liu H, Hu H (2018) Bacteria-mediated phthalic acid esters degradation and related molecular mechanisms. *Appl. Microbiol. Biotechnol*.
- Rodríguez-González V, Domínguez-Espíndola RB, Casas-Flores S, Patrón-Soberano OA, Camposeco-Solis R, Lee SW (2016) Antifungal nanocomposites inspired by titanate nanotubes for complete inactivation of *botrytis cinerea* isolated from tomato infection. *ACS Appl Mater Interfaces* 8:31625–31637 . <https://doi.org/10.1021/acsami.6b10060>
- Sandy EH, Yao J, Zheng S, Gogra AB, Chen H, Zheng H, Yormah TBR, Zhang X, Zaray G, Ceccanti B, Choi MMF (2010) A comparative cytotoxicity study of isomeric alkylphthalates to metabolically variant bacteria. *J Hazard Mater* 182:631–639 . <https://doi.org/10.1016/j.jhazmat.2010.06.079>
- Sharma N, Kumar V, Maitra SS, Lakkaboyana SK, Khantong S (2021) DBP biodegradation kinetics by *Acinetobacter* sp.33F in pristine agricultural soil. *Environ Technol Innov* 21:101240 . <https://doi.org/10.1016/j.eti.2020.101240>
- Singh N, Dalal V, Mahto JK, Kumar P (2017) Biodegradation of phthalic acid esters (PAEs) and in silico structural characterization of mono-2-ethylhexyl phthalate (MEHP) hydrolase on the basis of close structural homolog. *J Hazard Mater* 338:11–22 . <https://doi.org/10.1016/j.jhazmat.2017.04.055>
- Tang W, Zhang L, Fang Y, Zhou Y, Ye B (2016) Biodegradation of phthalate esters by newly isolated *Rhizobium* sp . LMB-1 and its biochemical pathway of di- n - butyl phthalate. 177–186 . <https://doi.org/10.1111/jam.13123>

- Tao Y, Li H, Gu J, Shi H, Han S, Jiao Y, Zhong G, Zhang Q, Akindolie MS, Lin Y, Chen Z, Zhang Y (2019) Ecotoxicology and Environmental Safety Metabolism of diethyl phthalate (DEP) and identification of degradation intermediates by *Pseudomonas* sp . DNE-S1. *Ecotoxicol Environ Saf* 173:411–418 . <https://doi.org/10.1016/j.ecoenv.2019.02.055>
- Wang P, Gao J, Zhao Y, Zhang M, Zhou S (2021a) Biodegradability of di-(2-ethylhexyl) phthalate by a newly isolated bacterium *Achromobacter* sp. RX. *Sci Total Environ* 755: . <https://doi.org/10.1016/j.scitotenv.2020.142476>
- Wang P, Gao J, Zhao Y, Zhang M, Zhou S (2021b) Biodegradability of di-(2-ethylhexyl) phthalate by a newly isolated bacterium *Achromobacter* sp. RX. *Sci Total Environ* 755:142476 . <https://doi.org/10.1016/j.scitotenv.2020.142476>
- Wang Y, Li F, Ruan X, Song J, Lv L, Chai L, Yang Z, Luo L (2017) Biodegradation of di-n-butyl phthalate by bacterial consortium LV-1 enriched from river sludge. *PLoS One*. <https://doi.org/10.1371/journal.pone.0178213>
- Wang Y, Miao B, Hou D, Wu X, Peng B (2012) Biodegradation of di-n-butyl phthalate and expression of the 3,4-phthalate dioxygenase gene in *Arthrobacter* sp. ZH2strain. *Process Biochem* 47:936–940 . <https://doi.org/10.1016/j.procbio.2012.02.027>
- Wang Y, Tam NFY (2018) *Microbial remediation of organic pollutants*, Second Edi. Elsevier Ltd.
- Wang Y, Yin B, Hong Y, Yan Y, Gu JD (2008) Degradation of dimethyl carboxylic phthalate ester by *Burkholderia cepacia* DA2 isolated from marine sediment of South China Sea. *Ecotoxicology* 17:845–852 . <https://doi.org/10.1007/s10646-008-0247-4>
- Wang Y, Zhu H, Kannan K (2019a) A review of biomonitoring of phthalate exposures. *Toxics* 7:1–28 . <https://doi.org/10.3390/TOXICS7020021>
- Wang Z, Zhu X, Su Y, Xu W, Liu H, Liu Z, Chen W, Wang J (2019b) Dimethyl phthalate damaged the cell membrane of *Escherichia coli* K12. *Ecotoxicol Environ Saf* 180:208–214 . <https://doi.org/10.1016/j.ecoenv.2019.05.009>
- Wittassek M, Koch HM, Angerer J, Brüning T (2011) Assessing exposure to phthalates - The human biomonitoring approach. *Mol Nutr Food Res* 55:7–31 .

<https://doi.org/10.1002/mnfr.201000121>

- Wu Q, Liu H, Ye LS, Li P, Wang YH (2013) Biodegradation of Di-n-butyl phthalate esters by *Bacillus* sp. SASHJ under simulated shallow aquifer condition. *Int Biodeterior Biodegrad* 76:102–107 . <https://doi.org/10.1016/j.ibiod.2012.06.013>
- Wu X, Liang R, Dai Q, Jin D, Wang Y, Chao W (2010) Complete degradation of di-n-octyl phthalate by biochemical cooperation between *Gordonia* sp. strain JDC-2 and *Arthrobacter* sp. strain JDC-32 isolated from activated sludge. *J Hazard Mater* 176:262–268 . <https://doi.org/10.1016/j.jhazmat.2009.11.022>
- Xu J, Lu Q, de Toledo RA, Shim H (2017) Degradation of di-2-ethylhexyl phthalate (DEHP) by an indigenous isolate *Acinetobacter* sp. SN13. *Int Biodeterior Biodegrad* 117:205–214 . <https://doi.org/10.1016/j.ibiod.2017.01.004>
- Xu XR, Li H Bin, Gu JD (2005) Biodegradation of an endocrine-disrupting chemical di-n-butyl phthalate ester by *Pseudomonas fluorescens* B-1. *Int Biodeterior Biodegrad* 55:9–15 . <https://doi.org/10.1016/j.ibiod.2004.05.005>
- Xu Y, Liu X, Zhao J, Huang H, Wu M, Li X, Li W, Sun X, Sun B (2021) An efficient phthalate ester-degrading *Bacillus subtilis*: Degradation kinetics, metabolic pathway, and catalytic mechanism of the key enzyme. *Environ Pollut* 273:116461 . <https://doi.org/10.1016/j.envpol.2021.116461>
- Zhang J, Zhang C, Zhu Y, Li J, Li X (2018) Biodegradation of seven phthalate esters by *Bacillus mojavensis* B1811. *Int Biodeterior Biodegrad*. <https://doi.org/10.1016/j.ibiod.2018.04.006>
- Zhang Y, Jiao Y, Li Z, Tao Y, Yang Y (2021) Hazards of phthalates (PAEs) exposure: A review of aquatic animal toxicology studies. *Sci. Total Environ*. 771:145418
- Zhao H, Hu R, Chen X, Chen X, Lü H, Li Y, Li H, Mo C, Cai Q, Wong M (2018a) Science of the Total Environment Biodegradation pathway of di- (2-ethylhexyl) phthalate by a novel *Rhodococcus pyridinivorans* XB and its bioaugmentation for remediation of DEHP contaminated soil. *Sci Total Environ* 640–641:1121–1131 . <https://doi.org/10.1016/j.scitotenv.2018.05.334>
- Zhao HM, Du H, Huang CQ, Li S, Zeng XH, Huang XJ, Xiang L, Li H, Li YW, Cai QY, Mo CH, He Z (2019) Bioaugmentation of Exogenous Strain *Rhodococcus*

- sp. 2G Can Efficiently Mitigate Di(2-ethylhexyl) Phthalate Contamination to Vegetable Cultivation. *J Agric Food Chem* 67:6940–6949 .
<https://doi.org/10.1021/acs.jafc.9b01875>
- Zhao HM, Du H, Lin J, Chen X Bin, Li YW, Li H, Cai QY, Mo CH, Qin HM, Wong MH (2016) Complete degradation of the endocrine disruptor di-(2-ethylhexyl) phthalate by a novel *Agromyces* sp. MT-O strain and its application to bioremediation of contaminated soil. *Sci Total Environ* 562:170–178 .
<https://doi.org/10.1016/j.scitotenv.2016.03.171>
- Zhao HM, Hu RW, Du H, Xin XP, Li YW, Li H, Cai QY, Mo CH, Liu JS, Zhou DM, Wong MH, He ZL (2018b) Functional genomic analysis of phthalate acid ester (PAE) catabolism genes in the versatile PAE-mineralising bacterium *Rhodococcus* sp. 2G. *Sci Total Environ* 640–641:646–652 .
<https://doi.org/10.1016/j.scitotenv.2018.05.337>
- Zhu F, Zhu C, Doyle E, Liu H, Zhou D, Gao J (2018) Fate of di (2-ethylhexyl) phthalate in different soils and associated bacterial community changes. *Sci Total Environ* 637–638:460–469 .
<https://doi.org/10.1016/j.scitotenv.2018.05.055>

8 Concluding remarks

The SSB-consortium is mainly composed by the genera *Serratia*, *Methylobacillus*, *Achromobacter*, and *Pseudomonas*.

The SSB-consortium degrades up to 500 mg L⁻¹ of DINP following a first-order kinetics.

The temperature and the initial concentration of DINP are the main factors affecting its biodegradation without accumulating toxic by-products.

De-esterification and β -oxidation are proposed as major biodegradation pathways based on the analysis of degradation intermediates.

These results indicate that the SSB-consortium is a potential alternative for environmental DINP removal.

9 Perspectives

The results obtained in this research showed that the SSB consortium is a potential candidate for bioremediation of DINP contaminated sites, however, a more complete understanding of their mechanisms and interactions could provide a basis for the development of future bioremediation strategies. Therefore, it would be interesting to investigate the following:

- To establish the microbial dynamics of the SSB-consortium during the biodegradation process.
- To evaluate the enzymatic activity of the SSB-consortium for the degradation of phthalates.
- To identify the genes involved in phthalate ester biodegradation pathways.
- To explore the effect of biodegradation of phthalates in different matrices such as soil and industrial effluents, to validate their degradation efficiency.


10 Annexes

Scientific articles

Paper 1. Pereyra-Camacho, M.A., Balderas-Hernández, V.E. & De Leon-Rodriguez, A. Biodegradation of diisononyl phthalate by a consortium of saline soil bacteria: optimisation and kinetic characterisation. *Appl Microbiol Biotechnol* 105, 3369–3380 (2021). <https://doi.org/10.1007/s00253-021-11255-5>

Paper 2. González-Escobar, J.L., **Pereyra-Camacho, M.A.**, De León-Rodríguez, A. et al. Biodegradation of recalcitrant compounds and phthalates by culturable bacteria isolated from *Liometopum apiculatum* microbiota. *World J Microbiol Biotechnol* 36, 73 (2020). <https://doi.org/10.1007/s11274-020-02850-1>

Congress conferences



Marco A. Pereyra-Camacho, Víctor E. Balderas-Hernández, Antonio De León-Rodríguez (2018). Bacterial biodegradation of endocrine disruptors di-(2-ethylhexyl) phthalate and di-isononyl phthalate. XXXII National Congress of Biochemistry. Ixtapa Zihuatanejo, Gro. Mexico.

Science communication conferences

LA EMPRESA LEBENS - CAPACITACIONES CIA. LTDA. CON EL AVAL ACADÉMICO DEL
INSTITUTO DE FORMACIÓN EMPRESARIAL AZTECA - ESTUDIOS UNIVERSITARIOS
(CENTRO DE NEXO ECUADOR)

 Lebens
OTORGAN EL PRESENTE
CERTIFICADO A:
Ing. Marco Antonio Pereyra M.S.C.

Por haber participado como **PONENTE** en el "III Congreso Internacional de Agroindustria: Tendencias Industriales, Biotecnología y Emprendimiento 2020", con el tema: **Biodegradación xenobióticos in agricultural soils; desarrollado los días 27, 28, 29 de noviembre y 04, 05, 06 de diciembre de 2020**, con una duración de 60 horas académicas.

Riobamba – Ecuador, diciembre de 2020.


Dr. Lesly Vanessa Salas
RECTORA
INSTITUTO DE FORMACIÓN EMPRESARIAL
AZTECA - ESTUDIOS UNIVERSITARIOS


Jorge Riquenne Domínguez
PRESIDENTE
ASOCIACIÓN NACIONAL DE ESTUDIANTES
AGROINDUSTRIALES DEL ECUADOR


Ing. Erick Escobar Ureña
PRESIDENTE
LEBENS - CAPACITACIONES CIA. LTDA.

 ANEA

 ANEA

 INSTITUTO
AZTECA
DE FORMACIÓN EMPRESARIAL

 III
CATIBE

Marco A. Pereyra-Camacho (2020). Biodegradation - Xenobiotics in Agricultural Soils. III International Congress on Agroindustry: Industrial Trends, Biotechnology and Entrepreneurship. Lebens. Riobamba, Ecuador.



Otorga a:

MSc. Marco Pereyra Camacho

el presente

Diploma

En el 9no Congreso de Biotecnología Quorum

Con su Conferencia Magistral

***“Biorremediación 3.0 - La Revolución de la
Biología en sistemas.”***

Querétaro, Qro. a 02 de octubre de 2020

Dra Grisel Fierros Romero
*Directora de Ingeniería en Biotecnología
Tecnológico de Monterrey Campus Querétaro*

Joana Lizeth Hernández Lara
*Presidente del 9no. Congreso Quorum
Tecnológico de Monterrey Campus Querétaro*

Marco A. Pereyra-Camacho (2020). Bioremediation 3.0. The Revolution of Systems Biology. 9th Edition of the International Biotechnology Congress QUORUM. Instituto Tecnológico y de Estudios Superiores de Monterrey, Campus Querétaro. Querétaro, Mexico.



El Instituto Tecnológico de Sonora

A través del Departamento de Biotecnología y Ciencias Alimentarias, el Programa Educativo de Ingeniero Biotecnólogo y la Asociación de Alumnos de Ingeniero Biotecnólogo



Otorga la presente

Constancia

a:


Dr. Marco A. Pereyra Camacho

Por su valiosa participación como Ponente en la "7ma Semana de Biotecnología" el día 12 de Marzo de 2019, con la conferencia "Ftalatos: amenazas silenciosas" en el Instituto Tecnológico de Sonora.



Dra. Olga Nydia Campas Baypoli

Jefe del Departamento de Biotecnología y Ciencias Alimentarias.



M.C. Eunice Guzmán Fierros

Responsable del Programa Educativo de Ingeniero Biotecnólogo



Juan Alberto Corral Palatox

Presidente de la Asociación de Alumnos de Ingeniero Biotecnólogo.

Cd. Obregón, Sonora, Marzo 2019.

Marco A. Pereyra-Camacho (2019). Phthalates: a silent threat. 7th Biotechnology Week of Sonora Institute of Technology (ITSON). Cd. Obregon, Mexico.