

This is the Author's Pre-print version of the following article: *Haret C. Rosu, Stefan C. Mancas, Chun-Chung Hsieh, Superfluid Rayleigh–Plesset extension of FLRW cosmology, Annals of Physics, Volume 429, 2021, 168490*, which has been published in final form at: [10.1016/j.aop.2021.168490](https://doi.org/10.1016/j.aop.2021.168490)

© 2021 This manuscript version is made available under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) license <http://creativecommons.org/licenses/by-nc-nd/4.0/>

# Superfluid Rayleigh-Plesset extension of FLRW cosmology

Haret C. Rosu\*, Stefan C. Mancas $\diamond$ , Chun-Chung Hsieh\*

\* *IPICYT- Instituto Potosino de Investigación Científica y Tecnológica,*

*Camino a la Presa San José 2055, Col. Lomas 4a Sección, San Luis Potosí, 78216 S.L.P., Mexico*

$\diamond$  *Department of Mathematics, Embry-Riddle Aeronautical University, Daytona Beach, FL 32114-3900, USA*

\* *Institute of Mathematics, Academia Sinica, Nankang, Taipei 115, Taiwan*

(Dated: 30 January 2021)

Guided by the analogy with the Rayleigh-Plesset dynamics of multielectron bubbles in superfluid He-4, we consider the cosmological FLRW evolution equation with additional cubic and sixth powers of the inverse of the scale factor of the universe. For zero cosmological constant along with the absence of viscous terms, and using the Sundman time as parameter, we present parametric solutions for the scale factor of the universe in terms of rational expressions of Weierstrass elliptic functions and special and degenerate cases thereof.

## I. INTRODUCTION

Hydrodynamic phenomena have long served as rich-in-wisdom analogies for astrophysics and cosmology. The Mach shock wave context of the black hole evaporation introduced by Unruh [1] four decades ago is just a famous example which attracted a lot of attention, but clearly it is not the only case. Superfluid hydrodynamics with its embodiments as Bose-Einstein condensates and superlight bosons is another notorious example in strong association with galactic halos and the concept of dark matter [2–4], whereas various other superfluid analogs of quantum field cosmic phenomena can be found in a review by Volovik [5]. Let us also recall that the idea of a superfluid universe has been central in the works of the late Professor Kerson Huang who summarized the publications with his collaborators along many years in an inspiring book on this topic [6].

On the other hand, in the latter case of cosmology, the fluid and classical mechanics analogies start to abound [7–10]. Among the fluid analogies, a less known, but equally promising one has been recently introduced by Rousseaux and Mancas [11] that we address here from the superfluid point of view. It is based on the similarities between the Rayleigh-Plesset (RP) equation [12, 13], which is a reshaped form of the Navier-Stokes equations describing the dynamical evolution of bubbles in fluids [14], and the cosmological dynamics as introduced by Friedmann, Lemaitre, Robertson, and Walker almost one century ago. The modern compelling components of the universe, such as dark energy and dark matter, may require the addition of supplementary terms to the latter equations. This is what Banerjee et al [15] have recently done by adding a fourth power term in the inverse of the scale factor as due to dark energy to enhance the late expansion rate of the universe, while Rousseaux and Mancas [11] speculated on the possible addition of more viscosity terms in the cosmological evolution, motivated by the case of the RP equation for which the addition of this kind of terms is a common usage, as revealed by some references cited in [11].

In this paper, motivated by [11], we investigate in an explicit way the addition of polynomial nonlinearities to the usual FLRW evolution equation. In particular, we are interested not only in the common cubic power of the usual RP equation, but also in the effects of an additional nonlinear power of order six which occurs in the Rayleigh-Plesset framework developed in the study of multielectron bubbles (henceforce mebs) in superfluid Helium-4 laboratory physics [16–18]. Such a term takes into account the repulsive Coulomb interaction of the electrons forming a thin layer at the surface of the bubbles. By considering such a phenomenological detail in theoretical cosmology, we place ourselves from the beginning in a particular case of the class of inhomogeneous fluid models of the universe which are widely discussed in the literature, see e.g. [19] and references therein. However, we believe that the detail that we emphasize here may be important not only for the laboratory phenomenology, but also in cosmology and astrophysics.

We also remark that the addition of power nonlinearities is well known in other research areas. It is sufficient to recall that a quintic power is added to the cubic nonlinear Schrödinger equation to handle the rich phenomenology of soliton propagation in nonlinear waveguides, and the case of laboratory BECs, where the cubic-quintic Ginzburg-Landau equation has been for several decades the standard equation for modeling the BEC behavior. Differently from the case of the single cubic nonlinearity which is integrable by the inverse scattering method, the solutions of the cubic quintic cases can develop blow-ups, finite time singularities where the amplitude of solution reaches infinity in a finite time [20]. The FLRW cosmology is not far from these areas in the dynamical nonlinear perspective. The standard FLRW equations with physical content characterized by normal (not ‘super-negative’-dark [21, 22]) forms of equation of state are integrable, see e.g. [23] and the references to textbooks therein, but adding more power nonlinearities can generate future-in-time singularities.

The rest of the paper is focused on obtaining some representative cosmological scaling factors of this extended FLRW model for zero cosmological constant which are obtained as solutions of the Weierstrass elliptic equation. In section II, we establish the nonlinear differential equation corresponding to this generalized FLRW model which by

a change of variable is turned into a Weierstrass elliptic equation whose standard mathematical setup is recalled. In section III, we present some sets of scale factors of the universe as parametric solutions of this model expressed in terms of rational expressions of Weierstrass elliptic  $\wp$  functions including the special and degenerate cases [24]. Finally, section IV contains some further remarks on the results and conclusions.

## II. THE FLRW EQUATION WITH SUPERFLUID RAYLEIGH-PLESSET TERMS AND THE ELLIPTIC EQUATION

The standard RP equation for the time-evolving radius,  $R(t)$ , of a bubble in a fluid reads

$$\frac{\ddot{R}}{R} + \frac{3}{2} \left( \frac{\dot{R}}{R} \right)^2 + \frac{-\Delta P(t)}{\rho_l} \left( \frac{1}{R} \right)^2 + \frac{2\gamma}{\rho_l} \left( \frac{1}{R} \right)^3 + 4\nu \left( \frac{\dot{R}}{R^3} \right) = 0, \quad (1)$$

to which a sixth power term in  $1/R$  should be added in the case of mebs. The symbols in the coefficients are:  $\rho_l$  – the liquid density,  $\nu$  – the liquid kinematic viscosity,  $\gamma$  – the surface tension, and  $\Delta P(t) = P_{in}(t) - P_{out}(t)$  – the relative pressure drop at the bubble with respect to the pressure in the liquid at infinity, and the dot denotes the time derivative. For this equation without the viscosity parametric solutions in terms of rational expressions of Weierstrass  $\wp$  functions have been already obtained by Kudryashov and Sinelshchikov [25].

By comparison, Rousseaux and Mancas [11] have proposed a  $\chi$  generalization of the single FLRW equation obtained from the coupled FLRW ones

$$\frac{\ddot{a}}{a} + \chi \left( \frac{\dot{a}}{a} \right)^2 + \chi k c^2 \left( \frac{1}{a} \right)^2 - \frac{1+\chi}{3} \Lambda c^2 + \alpha \left( \frac{1}{a} \right)^3 + \beta \left( \frac{1}{a} \right)^4 + \delta \left( \frac{\dot{a}}{a^3} \right) = 0. \quad (2)$$

where  $\chi = (1 + 3w)/2$ ,  $w$  is the parameter entering the barotropic equation of state  $p = w\rho c^2$ ,  $\rho$  is the total energy density with the standard vacuum, radiation, and matter components,  $k$  is the spatial curvature index, and  $\Lambda$  is the cosmological constant. The  $\alpha$  and  $\delta$  coefficients are cosmological equivalents of the corresponding Rayleigh-Plesset coefficients, and  $\beta$  is the coefficient of the term generated by the scenario developed in [15]. The equivalence  $a \equiv R$  serves as the basis of the analogy.

In this article, we study the simplified case when the quartic term is not taken into account and also the viscosity term is discarded since in the superfluid regime the viscosity is negligible. In addition, we will also assume  $\Lambda = 0$ , but on the other hand, we will include the sixth-power meb term. With these assumptions (2) reads

$$\frac{\ddot{a}}{a} + \chi \left( \frac{\dot{a}}{a} \right)^2 + \chi k c^2 \left( \frac{1}{a} \right)^2 + \alpha \left( \frac{1}{a} \right)^3 - \eta \left( \frac{1}{a} \right)^6 = 0. \quad (3)$$

We consider the case  $\chi = 3/2$  which is in direct correspondence with the RP equation and represents the case of monoatomic gas. Using natural units for the speed of light,  $c = 1$ , we write (3) as

$$a\ddot{a} + \frac{3}{2}\dot{a}^2 = \eta a^{-4} - \alpha a^{-1} - \frac{3}{2}k. \quad (4)$$

Multiplying (4) by the integrating factor  $2a^2\dot{a}$  and integrating once, we obtain

$$\dot{a}^2 = -2\eta a^{-4} + 4c_1 a^{-3} - \alpha a^{-1} - k, \quad (5)$$

where  $c_1$  is an arbitrary constant of integration that is found from two initial conditions. For  $\dot{a}(0) = 0$  and  $a(0) = a_0$  then  $c_1 = c_1(\alpha, k, \eta) \equiv (2\eta + \alpha a_0^3 + k a_0^4)/4a_0$ . Notice that  $a_0$  is a root of the right hand side of (5) and also of the polynomial  $Q(a)$  in (6) below. We shall use the symbol  $\mu$  for  $a_0$  in the rest of the paper. Most of the solutions that will be displayed in this paper do not have the big bang initial condition  $a_0 = \mu = 0$ , but the initial Hubble parameter  $H(0) = \dot{a}(0)/\mu$  is zero and also one can tune  $c_1$  to get an appropriate  $\mu$ .

Using the Sundman transformation  $dt = a^2 d\tau$  we obtain the elliptic equation

$$a_\tau^2 = -ka^4 - \alpha a^3 + 4c_1 a - 2\eta \equiv Q(a), \quad (6)$$

where the  $\tau$  subindex denotes the  $\tau$  derivative. This equation can be viewed as an energy conservation equation for a one degree of freedom classical particle of mass  $m = 2$  if written in the form  $a_\tau^2 + V(a) = E$ , where the potential energy is  $V(a) = a(ka^3 + \alpha a^2 - 4c_1)$  and the total energy is  $E(a, \dot{a}) = -2\eta = a^4 \dot{a}^2 - 4c_1 a + \alpha a^3 + ka^4$ .

### A. The General Mathematical Setup of Elliptic Equations

It is well known [26–28] that the solutions  $a(\tau)$  of

$$a_\tau^2 = c_4 a^4 + 4c_3 a^3 + 6c_2 a^2 + 4c_1 a + c_0, \quad (7)$$

can be expressed in terms of Weierstrass elliptic functions  $\wp(\tau; g_2, g_3)$ , which is a solution to

$$\wp_\tau^2 = 4\wp^3 - g_2\wp - g_3, \quad (8)$$

via the transformation

$$a(\tau) = \hat{a} + \frac{\sqrt{Q(\hat{a})}\wp'(\tau - \tau_0; g_2, g_3) + \frac{1}{2}Q'(\hat{a})\left[\wp(\tau - \tau_0; g_2, g_3) - \frac{1}{24}Q''(\hat{a})\right] + \frac{1}{24}Q(\hat{a})Q^{(3)}(\hat{a})}{2\left[\wp(\tau - \tau_0; g_2, g_3) - \frac{1}{24}Q''(\hat{a})\right]^2 - \frac{1}{48}Q(\hat{a})Q^{(4)}(\hat{a})}, \quad (9)$$

where  $\hat{a}$  is not necessarily a root of  $Q(a)$ , and  $g_2, g_3$  are elliptic invariants of  $\wp(\tau)$ , which satisfy

$$\begin{aligned} g_2 &= c_4 c_0 - 4c_3 c_1 + 3c_2^2 = 2k\eta + c_1\alpha \\ g_3 &= c_4 c_2 c_0 + 2c_3 c_2 c_1 - c_4 c_1^2 - c_2^3 - c_3^2 c_0 = kc_1^2 + \frac{1}{8}\alpha^2\eta. \end{aligned} \quad (10)$$

These invariants, with the varying parameter  $c_1(\alpha, k, \eta)$  together with the modular discriminant

$$\Xi = g_2^3 - 27g_3^2 \quad (11)$$

are used to classify the solutions of (6). In particular, choosing  $\mu$  to be any real root of  $Q(a)$ , i.e.  $Q(\mu) = 0$ , then the general solution (9) takes the much simpler form

$$a(\tau) = \mu + \frac{Q_a(\mu)}{4\wp(\tau - \tau_0; g_2; g_3) - \frac{Q_{aa}(\mu)}{6}}. \quad (12)$$

This solution can be also interpreted as the elliptic modulation,  $M_Q(\wp; c_1, \mu)$ , of any solution of the polynomial  $Q(a)$  if we write it in the form

$$a(\tau) = \mu \left( 1 + \frac{\frac{Q_a(\mu)}{\mu}}{4\wp(\tau - \tau_0; g_2; g_3) - \frac{Q_{aa}(\mu)}{6}} \right) \equiv \mu M_Q(\wp; c_1, \mu)$$

and can be generalized using (9) to any point  $\hat{a}, \hat{\tau}$  of the plane  $(a, \tau)$ .

## III. THE PARAMETRIC SOLUTIONS

We are now ready to present a whole wealth of parametric solutions of this extended FLRW model using  $\tau$  as parameter. Although all possible cases can be enlisted in a systematic manner, for a superfluid universe or one containing superfluid bubbles one should consider the simultaneous presence of the Laplace pressure and electrostatic terms, i.e., both  $\alpha$  and  $\eta$  should not be zero simultaneously. When both of them are chosen to be zero it corresponds to the standard FLRW cosmology. The case when one of them is zero such as  $\alpha = 0$  and  $\eta \neq 0$  are not physical, while  $\alpha \neq 0$  and  $\eta = 0$  describes common liquids, and thus will not be of interest here.

### A. Curved Space Cases, $k \neq 0, \Xi \neq 0$

Substituting  $Q(\mu)$  and its derivatives, (12) becomes the parametric *rational Weierstrass* solution

$$a(\tau) = \mu + \frac{4c_1 - 4k\mu^3 - 3\alpha\mu^2}{4\wp(\tau - \tau_0; 2k\eta + \alpha c_1; kc_1^2 + \frac{1}{8}\alpha^2\eta) + 2k\mu^2 + \alpha\mu}, \quad \alpha \neq 0, \quad \eta \neq 0. \quad (13)$$

Next, we choose two particular cases of nondegenerate solutions, one referred as superfluid rational lemniscatic for  $g_3 = 0$ , and the other as standard rational equianharmonic for  $g_2 = 0$  because it corresponds to the standard FLRW cosmology for which  $\alpha = \eta = 0$ .

i) The lemniscatic case: For  $c_1 = \pm \frac{\alpha}{4} \sqrt{-\frac{2\eta}{k}}$  obtained by solving  $g_3 = 0$ , and  $\mu$  a root of  $-ka^4 - \alpha a^3 \pm \alpha \sqrt{-\frac{2\eta}{k}} a - 2\eta = 0$  we obtain the rational *lemniscatic* solution

$$a(\tau) = \mu + \frac{\pm \alpha \sqrt{-\frac{2\eta}{k}} - 4k\mu^3 - 3\alpha\mu^2}{4\wp\left(\tau - \tau_0; 2k\eta \pm \frac{\alpha^2}{4} \sqrt{-\frac{2\eta}{k}}; 0\right) + 2k\mu^2 + \alpha\mu}, \quad \alpha \neq 0, \quad \eta \neq 0, \quad g_3 = 0. \quad (14)$$

Plots of the periodic (closed universe) and non periodic (open universe) lemniscatic cases are displayed in Figs. 1 and 2, respectively. One can see the very interesting feature that the physical solutions of the periodic cases have contributions from the initially negative solutions.




FIG. 1: Rational lemniscatic solutions for a superfluid FLRW closed universe from (14).

ii) Standard FLRW cosmology,  $\alpha = 0$  and  $\eta = 0$ : Let  $\mu$  be a root of  $-ka^4 + 4c_1a = 0$ , then (13) becomes the *equianharmonic* solution

$$a(\tau) = \mu + \frac{2(c_1 - k\mu^3)}{2\wp(\tau - \tau_0; 0; kc_1^2) + k\mu^2}, \quad \alpha = 0, \quad \eta = 0, \quad g_2 = 0. \quad (15)$$

The equianharmonic periodic case corresponding to the closed universe is plotted in Fig. 3, while the non periodic case of the open universe is displayed in Fig. 4. As can be seen in these figures, the solutions are in the two classes of either pure positive (physical) or pure negative (nonphysical).




FIG. 2: The same type of solutions for the superfluid open universe from (14).




FIG. 3: Rational equianharmonic solutions for a standard FLRW ( $\alpha = 0$ ,  $\eta = 0$ ) closed universe from (15).

### B. Degenerate Modular Cases, $k \neq 0$ , $\Xi = 0$

We now study the degenerate cases when  $\Xi = 0$  by finding numerically the coefficient  $c_1$  by solving  $\Xi = 0$ . In this case  $\wp(\eta, g_2, g_3)$  degenerates into *hyperbolic*, *trigonometric* functions, or *rational* functions, and (13) can be simplified further as follows:




FIG. 4: Same type of solutions for a standard FLRW ( $\alpha = 0$ ,  $\eta = 0$ ) open universe from (15).

i) Let  $g_2 = 12\hat{e}^2 > 0$  and  $g_3 = -8\hat{e}^3 < 0$ ; the Weierstrass  $\wp$  function is simplified to

$$\wp(\tau; 12\hat{e}^2, -8\hat{e}^3) = \hat{e} \left[ 1 + 3 \operatorname{csch}^2(\sqrt{3}\hat{e} \tau) \right]. \quad (16)$$

Using this in (13) we obtain the rational *hyperbolic* solutions

$$a(\tau) = \mu + \frac{4c_1^* - 4k\mu^3 - 3\alpha\mu^2}{2\sqrt[3]{-g_3} \left[ 1 + 3\operatorname{csch}^2\left(\frac{\sqrt{6}}{2}\sqrt[3]{-g_3}(\tau - \tau_0)\right) \right] + 2k\mu^2 + \alpha\mu}, \quad g_2 > 0, \quad g_3 < 0. \quad (17)$$

As shown in Fig. 5, these are solutions reaching quickly plateau regions for both types of curved spaces.




FIG. 5: Rational degenerate hyperbolic solutions for superfluid closed and open universe from (17), left and right, respectively.

ii) Let  $g_2 = 12\tilde{e}^2 > 0$  and  $g_3 = 8\tilde{e}^3 > 0$ ; the Weierstrass  $\wp$  function is simplified to

$$\wp(\tau; 12\tilde{e}^2, 8\tilde{e}^3) = \tilde{e} \left[ -1 + 3 \operatorname{csc}^2(\sqrt{3}\tilde{e} \tau) \right]. \quad (18)$$

Thus, we obtain the rational *trigonometric* solutions

$$a(\tau) = \mu + \frac{4c_1^* - 4k\mu^3 - 3\alpha\mu^2}{2\sqrt[3]{g_3} \left[ -1 + 3\operatorname{csc}^2\left(\frac{\sqrt{6}}{2}\sqrt[3]{g_3}(\tau - \tau_0)\right) \right] + 2k\mu^2 + \alpha\mu}, \quad g_2 > 0, \quad g_3 > 0. \quad (19)$$

As shown in the plots of Fig. 6, the closed universe case can have contributions from the initially unphysical trigonometric solution.




FIG. 6: Rational degenerate trigonometric solutions for superfluid closed and open universe from (19).

iii) When both germs are zero then  $\wp(\tau, 0, 0) = \frac{1}{\tau^2}$ , and we have the *rational* solution

$$a(\tau) = \mu + \frac{Q_a(\mu)\eta^2}{4 - \frac{Q_{aa}(\mu)}{6}\eta^2}, \quad (20)$$

which simplifies to

$$a(\tau) = \mu + \frac{(4c_1 - 4k\mu^3 - 3\alpha\mu^2)\tau^2}{4 + (2k\mu^2 + \alpha\mu)\tau^2}, \quad g_2 = g_3 = 0. \quad (21)$$

Moreover, this case leads to  $c_1 = \frac{\alpha^3}{16k^2}$ , and  $\eta = -\frac{\alpha^4}{32k^3}$ , thus the quartic polynomial has four real zeros inferred from  $Q(\mu) = \frac{1}{16k^3}(\alpha - 2k\mu)(\alpha + 2k\mu)^3$ , so the graph of (21) becomes the *Agnesi* curve [29]

$$a(\tau) = \frac{\alpha}{2} - \frac{\alpha^3\tau^2}{4 + \alpha^2\tau^2} = \frac{\alpha}{2} \left( 1 - \alpha\tau^2 \frac{d}{d\tau} \arctan(\alpha\tau/2) \right), \quad k = 1, \quad g_2 = g_3 = 0, \quad (22)$$

$$a(\tau) = -\frac{\alpha}{2} - \frac{\alpha^3\tau^2}{4 - \alpha^2\tau^2} = -\frac{\alpha}{2} \left( 1 + \alpha\tau^2 \frac{d}{d\tau} \operatorname{arctanh}(\alpha\tau/2) \right), \quad k = -1, \quad g_2 = g_3 = 0.$$

The degenerate Agnesi solutions in curved spaces are presented in Fig. 7, and again one can see that it is possible that the initially negative solution can become the physical solution after some time in the case of closed universes.




FIG. 7: Degenerate Agnesi solutions for superfluid closed and open universe from (22), left and right, respectively.

For all cases, the comoving time in parametric form is obtained from  $dt = a^2 d\tau$  which gives then

$$t(\tau) = \int_{\tau_0}^{\tau} a^2(\xi) d\xi. \quad (23)$$



### C. Flat Space Cases, $k = 0$ , $\Xi \neq 0$

In this case, (6) simplifies to the elliptic equation

$$a_\tau^2 = -\alpha a^3 + 4c_1 a - 2\eta \quad (24)$$

which can be reduced to the Weierstrass elliptic equation (8) by the scale transformation  $a(\tau) = -\frac{4}{\alpha}\wp(\tau - \tau_0; g_2, g_3)$ , where

$$\begin{aligned} g_2 &= c_1 \alpha, \\ g_3 &= \frac{1}{8} \alpha^2 \eta \\ \Xi &= (c_1 \alpha)^3 - \frac{27}{64} (\alpha^2 \eta)^2 \end{aligned} \quad (25)$$

and so, the parametric *Weierstrass* solutions for  $k = 0$  are

$$\begin{aligned} a(\tau) &= -\frac{4}{\alpha} \wp \left( \tau - \tau_0; c_1 \alpha, \frac{1}{8} \alpha^2 \eta \right), \quad \alpha \neq 0, \quad \eta \neq 0 \\ t(\tau) &= \int_{\tau_0}^{\tau} a^2(\xi) d\xi = \frac{4}{3\alpha^2} \left[ c_1 \alpha (\tau - \tau_0) + \hat{c}_1 + 2\wp' \left( \tau; c_1 \alpha, \frac{1}{8} \alpha^2 \eta \right) \right], \quad \tau_0 \neq 0, \end{aligned} \quad (26)$$

where  $\hat{c}_1 = 2\wp'(\tau_0; c_1 \alpha, \frac{1}{8} \alpha^2 \eta)$ . We will consider only the simpler equianharmonic subcase  $c_1 = 0$ , but still describing a superfluid FLRW (flat) universe.

i)  $c_1 = 0$ : one obtains the reduced elliptic equation

$$a_\tau^2 = -\alpha a^3 - 2\eta. \quad (27)$$

The solutions can be obtained directly from (26) to give

$$\begin{aligned} a(\tau) &= -\frac{4}{\alpha} \wp \left( \tau - \tau_0; 0, \frac{1}{8} \alpha^2 \eta \right), \quad \alpha \neq 0, \quad \eta \neq 0, \quad c_1 = 0, \\ t(\tau) &= \int_{\tau_0}^{\tau} a^2(\xi) d\xi = \frac{8}{3\alpha^2} \left[ \hat{c}_1 + \wp' \left( \tau; 0, \frac{1}{8} \alpha^2 \eta \right) \right], \quad \tau_0 \neq 0, \end{aligned} \quad (28)$$

where  $\hat{c}_1 = \wp'(\tau_0; 0, \frac{1}{8} \alpha^2 \eta)$ . For this reduced *equianharmonic* case it is also possible to have physical contributions from the initially negative solution as seen in Fig. 8.




FIG. 8: Reduced equianharmonic solutions for a superfluid FLRW flat space from (28) for  $\tau_0 = 1$  which avoids the singularity of  $\wp'$  at the origin.

### D. Degenerate Modular Cases, $k = 0$ , $\Xi = 0$

These cases are obtained for  $c_1^* = \frac{3}{4} \sqrt[3]{\alpha \eta^2}$  which leads to hyperbolic or trigonometric solutions, or when  $\alpha = 0$  which is physical as standard FLRW only when  $\eta = 0$ .

i) Let  $g_2 = 12\hat{e}^2 = \frac{3}{4}\alpha^{4/3}\eta^{2/3} > 0$  and  $g_3 = -8\hat{e}^3 = \frac{1}{8}\alpha^2\eta < 0$ . These values of the germs lead to the *hyperbolic* solutions

$$a(\tau) = -\frac{2}{\alpha} \sqrt[3]{-g_3} \left[ 1 + 3\text{csch}^2 \left( \frac{\sqrt{6}}{2} \sqrt[6]{-g_3} (\tau - \tau_0) \right) \right], \quad g_2 > 0, \quad g_3 < 0. \quad (29)$$

ii) Let  $g_2 = 12\hat{e}^2 = \frac{3}{4}\alpha^{4/3}\eta^{2/3} > 0$  and  $g_3 = 8\hat{e}^3 = \frac{1}{8}\alpha^2\eta > 0$ . For these germs, one obtains the *trigonometric* solutions

$$a(\tau) = -\frac{2}{\alpha} \sqrt[3]{g_3} \left[ -1 + 3\text{csc}^2 \left( \frac{\sqrt{6}}{2} \sqrt[6]{g_3} (\tau - \tau_0) \right) \right], \quad g_2 > 0, \quad g_3 > 0. \quad (30)$$

iii)  $\alpha = 0$  and  $\eta = 0$ : then (24) becomes

$$a_\tau^2 = 4c_1 a \quad (31)$$

with parametric *polynomial* solutions

$$\begin{aligned} a(\tau) &= \pm 2\sqrt{c_1}(\tau - \tau_0), \quad \alpha = 0, \quad \eta = 0 \\ t(\tau) &= \frac{4}{3}c_1\tau(\tau^2 - 3\tau\tau_0 + 3\tau_0^2). \end{aligned} \quad (32)$$

Examples of the degenerate hyperbolic, trigonometric, and the last class of polynomial solutions in flat space are illustrated in Fig. 9 without showing contributions from the initially negative solutions for the employed parameters.




FIG. 9: Modular degenerate solutions of hyperbolic, trigonometric, and polynomial type corresponding to the superfluid FLRW flat space.

#### IV. DISCUSSION AND CONCLUSION

We have introduced and analyzed in some detail a nonlinear differential equation for a modified FLRW cosmology for zero cosmological constant, which is based on an analogy with a Rayleigh-Plesset equation encountered in the area of multielectron bubbles in superfluid He-4, a well studied laboratory phenomenon. At the speculative level, one can assume the existence of equivalents of such bubbles even at astrophysical scales, as a kind of microcavities hovering in the galactic halos where they could be generated due to some unknown astrophysical mechanism.

We have presented parametric analytical solutions which are mainly in rational forms containing elliptic Weierstrass  $\wp$  functions in terms of a variable that we call Sundman time whose differential is  $a^2$  times smaller than the differential of the comoving time. For standard FLRW cosmology, some results in the comoving time involving the Weierstrass  $\wp$  function goes back to Lemaître himself [30], while Coquereaux [31] obtained an equianharmonic (nonrational)  $\wp$  solution in conformal time when the radiative contribution is neglected. The reduction to the standard FLRW case of our superfluid RP-modified FLRW cosmology provides the rational equianharmonic solution  $a(\tau) = c_1/\wp(\tau; 0, kc_1^2)$  for  $\mu = 0$  which exists only if  $c_1 \neq 0$ , see equation (15) and Figs 3(a) and 4(a). A similar result in conformal time can

be found in the work of Steiner [24], but with  $\wp$  not strictly equianharmonic. However, this author claims a correction of sign in the main formula of the first Biermann-Weierstrass theorem in [27] that we do not confirm.

For some particular values of the parameters, we have displayed periodic (cyclic) solutions specific to rational expressions of Weierstrass functions. The interesting feature of physical contributions from an initially nonphysical (negative) solution has been revealed, mainly but not only, for the closed universe cases. In the case in which some of these superfluid-modified scale factors will be taken into account for possible cosmological scenarios or for astrophysical phenomena, then the parameters should be fitted to more realistic values according to the sets of observational data. Furthermore, at the terrestrial laboratory level, some of the flat space solutions that have been obtained here can be also useful in the data analyses of superfluid bubble experiments.

- 
- [1] W.G. Unruh, Experimental black-hole evaporation?, *Phys. Rev. Lett.* **46**, 1351-1353 (1981).
- [2] S. J. Sin, Late time cosmological phase transition and galactic halo as Bose liquid, *Phys. Rev. D* **50**, 3650-3654 (1994).
- [3] X. Zhang, M.H. Chan, T. Harko, S.-D. Liang, and C.S. Leung, Slowly rotating BEC galactic dark matter halos, and their rotation curves, *Eur. Phys. J. C* **78**, 346 (2018).
- [4] L. Hui, J.P. Ostriker, S. Tremaine, and E. Witten, Ultralight scalars as cosmological dark matter, *Phys. Rev. D* **95**, 043541 (2017).
- [5] G.E. Volovik, Superfluid analogies of cosmological phenomena, *Phys.Rept.* **351**, 195-348 (2001).
- [6] K. Huang, *A Superfluid Universe* (World Scientific, Singapore, 2016).
- [7] D. Bini and S. Succi, Analogy between capillary motion and Friedmann-Robertson-Walker cosmology, *Europhys. Lett.* **82**, 34003 (2008).
- [8] V. Faraoni, Analogy between freezing lakes and the cosmic radiation era, *Phys. Rev. Res.* **2**, 013187 (2020).
- [9] V. Faraoni, Cosmic analogues of classic variational problems, *Universe* **2020**, 6, 71 (2020).
- [10] M.N. Shneider and M. Pekker, Cavitation model of the inflationary stage of Big Bang, *Phys. Fluids* **33**, 017116 (2021).
- [11] G. Rousseaux and S.C. Mancas, Visco-elastic cosmology for a sparkling universe?, *Gen. Rel. Grav.* **52**, 55 (2020) [arXiv:2002.12123].
- [12] Lord Rayleigh, The pressure developed in a liquid during the collapse of a spherical cavity, *Phil. Mag.* **34**, 94-98 (1917).
- [13] M.S. Plesset, The dynamics of cavitation bubbles, *J. Appl. Mech.* **16**, 277-282 (1949).
- [14] R.A. Van Gorder, Dynamics of the Rayleigh-Plesset equation modelling a gas-filled bubble immersed in an incompressible fluid, *J. Fluid. Mech.* **807**, 479-508 (2016).
- [15] S. Banerjee, U. Danielsson, G. Dibitetto, S. Giri, and M. Schillo, Emergent de Sitter cosmology from decaying Anti-de Sitter space, *Phys. Rev. Lett.* **121**, 261301 (2018).
- [16] M.M. Salomaa and G.A. Williams, Structure and stability of multielectron bubbles in liquid Helium, *Phys. Rev. Lett.* **47**, 1730-1733 (1981).
- [17] J. Tampere, I.F. Silvera, and J.T. Devreese, Multielectron bubbles in helium as a paradigm for studying electrons on surfaces with curvature, *Surf. Sci. Rep.* **62**, 159-217 (2007).
- [18] N. Yadav, P.K. Rath, Z. Xie, Y. Huang, and A. Gosh, Multielectron bubbles in liquid Helium, *J. Low Temp. Phys.* **200**, xxx (2020).
- [19] S. Nojiri and S.D. Odintsov, Inhomogeneous equation of state of the universe: Phantom era, future singularity, and crossing the phantom barrier, *Phys. Rev. D* **72**, 023003 (2005).
- [20] I. S Aranson and L. Kramer, The world of the complex Ginzburg-Landau equation, *Rev. Mod. Phys.* **74**, 99-143 (2001).
- [21] S. Nojiri and S.D. Odintsov, The dark universe future and singularities: the account of thermal and quantum effects, *Physics of the Dark Universe* **30**, 100695 (2020) [arXiv:2006.03946].
- [22] R.R. Caldwell, A phantom menace ? Cosmological consequences of a dark energy component with super-negative equation of state, *Phys. Lett. B* **545**, 23-29 (2002).
- [23] V. Faraoni, Solving for the dynamics of the universe, *Am. J. Phys.* **67**, 732-734 (1999).
- [24] For the standard FLRW model, a discussion in terms of Weierstrass elliptic  $\wp$  functions can be found in F. Steiner, Solution of the Friedmann equation determining the time evolution, acceleration, and the age of the universe, [https://www.uniulm.de/fileadmin/website\\_uni\\_ulm/nawi.inst.260/paper/08/tp08-7.pdf](https://www.uniulm.de/fileadmin/website_uni_ulm/nawi.inst.260/paper/08/tp08-7.pdf), 1-18 (2007).
- [25] N.A. Kudryashov and D.I. Sinelshchikov, Analytical solutions for problems of bubble dynamics, *Phys. Lett. A* **379**, 798-802 (2015).
- [26] K. Weierstrass, *Mathematische Werke, vol. V* (Johnson, New York, 1915).
- [27] E.T. Whittaker and G.N. Watson, *A Course of Modern Analysis* (Cambridge University Press, Cambridge, 1927).
- [28] M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions: With Formulas, Graphs, and Mathematical Tables* (Courier Dover Publications, New York, 1972).
- [29] M.G. Agnesi, *Istituzioni analitiche ad uso della gioventú italiana*. See in particular Problem 3, 380-382, and Fig. 135 (1748).
- [30] G. Lemaitre, L'univers en expansion, *Ann. Soc. Sci. Brux. A* **53**, 51-85 (1933); English translation: The expanding universe, *Gen. Rel. Grav.* **29**, 641-80 (1997).
- [31] R. Coquereaux, The history of the universe is an elliptic curve, *Class. Quant. Grav.* **32**, 115013 (2015).