

This document is protected by copyright and was first published by Frontiers Media S.A. All rights reserved. It is reproduced with permission.

Citation:

Cuellar-Santoyo AO, Ruiz-Rodríguez VM, Mares-Barbosa TB, Patrón-Soberano A, Howe AG, Portales-Pérez DP, Miquelajáuregui Graf A and Estrada-Sánchez AM (2023) Revealing the contribution of astrocytes to glutamatergic neuronal transmission. *Front. Cell. Neurosci.* 16:1037641. doi: [10.3389/fncel.2022.1037641](https://doi.org/10.3389/fncel.2022.1037641)

Copyright © 2023 Cuellar-Santoyo, Ruiz-Rodríguez, Mares-Barbosa, Patrón-Soberano, Howe, Portales-Pérez, Miquelajáuregui Graf and Estrada-Sánchez. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



OPEN ACCESS

EDITED BY

Daniel Reyes-Haro,
Institute of Neurobiology, National
Autonomous University of Mexico, Mexico

REVIEWED BY

Sergei Kirischuk,
Johannes Gutenberg University Mainz,
Germany
Friederike Klempin,
Charité Universitätsmedizin Berlin, Germany

*CORRESPONDENCE

Ana María Estrada-Sánchez
✉ ana.estrada@ipicyt.edu.mx

SPECIALTY SECTION

This article was submitted to
Non-Neuronal Cells,
a section of the journal
Frontiers in Cellular Neuroscience

RECEIVED 06 September 2022

ACCEPTED 20 December 2022

PUBLISHED 19 January 2023

CITATION

Cuellar-Santoyo AO, Ruiz-Rodríguez VM,
Mares-Barbosa TB, Patrón-Soberano A,
Howe AG, Portales-Pérez DP, Miquelajáuregui
Graf A and Estrada-Sánchez AM
(2023) Revealing the contribution of astrocytes
to glutamatergic neuronal transmission.
Front. Cell. Neurosci. 16:1037641.
doi: 10.3389/fncel.2022.1037641

COPYRIGHT

© 2023 Cuellar-Santoyo, Ruiz-Rodríguez,
Mares-Barbosa, Patrón-Soberano, Howe,
Portales-Pérez, Miquelajáuregui Graf and
Estrada-Sánchez. This is an open-access article
distributed under the terms of the [Creative
Commons Attribution License \(CC BY\)](#). The
use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in this
journal is cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Revealing the contribution of astrocytes to glutamatergic neuronal transmission

Ares Orlando Cuellar-Santoyo¹, Victor Manuel Ruiz-Rodríguez¹,
Teresa Belem Mares-Barbosa^{1,2}, Araceli Patrón-Soberano¹,
Andrew G. Howe³, Diana Patricia Portales-Pérez²,
Amaya Miquelajáuregui Graf⁴ and Ana María Estrada-Sánchez^{1*}

¹División de Biología Molecular, Laboratorio de Neurobiología, Instituto Potosino de Investigación Científica y Tecnológica (IPICYT), San Luis Potosí, Mexico, ²Translational and Molecular Medicine Laboratory, Research Center for Health Sciences and Biomedicine, Autonomous University of San Luis Potosí, San Luis Potosí, Mexico, ³Intelligent Systems Laboratory, HRL Laboratories, LLC, Malibu, CA, United States, ⁴Faculty of Medicine and Health Sciences, University of Antwerp, Antwerp, Belgium

Research on glutamatergic neurotransmission has focused mainly on the function of presynaptic and postsynaptic neurons, leaving astrocytes with a secondary role only to ensure successful neurotransmission. However, recent evidence indicates that astrocytes contribute actively and even regulate neuronal transmission at different levels. This review establishes a framework by comparing glutamatergic components between neurons and astrocytes to examine how astrocytes modulate or otherwise influence neuronal transmission. We have included the most recent findings about the role of astrocytes in neurotransmission, allowing us to understand the complex network of neuron-astrocyte interactions. However, despite the knowledge of synaptic modulation by astrocytes, their contribution to specific physiological and pathological conditions remains to be elucidated. A full understanding of the astrocyte's role in neuronal processing could open fruitful new frontiers in the development of therapeutic applications.

KEYWORDS

calcium, gliotransmission, NMDA receptors, GLT-1, GLAST, VGLUT1, xCT

Introduction

The brain is one of the most complex organs in the human body; it consists of many distinct cell types, but most cells fall under the broad categories of neurons or glia with approximately even numbers in each category. The glial cells are divided into microglia, oligodendrocytes, and astrocytes; the latter comprise around 20% of the cells in the brain (Ventura and Harris, 1999; Salas et al., 2020). Astrocytes participate in many neurophysiological processes, including synaptogenesis (Allen and Eroglu, 2017), modulation of synaptic transmission, neuronal plasticity (Newman and Zahs, 1998; Araque et al., 1999), and regulation of blood flow in addition to the trafficking of small molecules and ions through their end-feet processes at the blood-brain barrier (Giaume et al., 1997; Simard et al., 2003). During physiological conditions, neurons and astrocytes have a coordinated functional relationship that ensures proper information flow, and each contributes to synaptic transmission by releasing neurotransmitters (by presynaptic neurons) or gliotransmitters (by astrocytes; see below).

Glutamate is the primary excitatory neurotransmitter in the mammalian brain, and it participates in diverse physiological processes such as learning, memory, and neuronal development (Yu et al., 1984; Behar et al., 1999; Hrabetova et al., 2000). However, glutamate can induce neuronal damage through excitotoxicity, which results from the over-activation of

glutamatergic receptors. Glutamate-mediated toxicity has been implicated in the pathogenesis of neurodegenerative diseases including Alzheimer's, Huntington's, and Parkinson's diseases (Koutsilieri and Riederer, 2007; Estrada-Sánchez et al., 2009; Ong et al., 2013), as well as psychiatric disorders such as schizophrenia (O'Donovan et al., 2017; Shah et al., 2020). Although glutamate toxicity contributes to these neuropathological conditions, the changes that lead to impaired glutamatergic neurotransmission are diverse, with different causes in each pathology. Nonetheless, an impaired relationship between neurons and astrocytes might be a common component. In the next section, we review the similarities and differences among the components of glutamatergic transmission in neurons and astrocytes.

Glutamatergic neurotransmission

Glutamate synthesis

Glutamate plays a role in multiple biological processes in the brain, and yet it cannot cross the blood-brain barrier. Instead, the brain's glutamate is synthesized locally by astrocytes and neurons through one of several pathways. We provide a brief description of glutamate metabolism (this section and Figure 1; Hawkins, 2009; Fernstrom, 2018); further details can be found in the extensive review by Schousboe et al. (2014).

The main precursor of glutamate in the brain is glutamine, a key component of the glutamine-glutamate cycle that encompasses the exchange of glutamine and glutamate between astrocytes and neurons. The reuptake of glutamate by its transporters, glutamate transporter 1 (GLT-1) and glutamate aspartate transporter (GLAST) in astrocytes enables the synthesis of glutamine by the cytosolic enzyme glutamine synthetase, which is highly expressed in astrocytes. Glutamine synthetase promotes the conversion of glutamate to glutamine using the cofactors NADPH, ATP, and NH_4^+ (Lehre et al., 1995; Bergles and Jahr, 1997; Anlauf and Derouiche, 2013; Huyghe et al., 2014; Yamada et al., 2019). Glutamine is then transported to the synaptic cleft, where neurons take it up through specific glutamine transporters (i.e., members of the families SNAT, LAT, ASC, and $^+$ LAT; Anlauf and Derouiche, 2013; Yamada et al., 2019). Once transported into neurons, the mitochondrial enzyme glutaminase (an amidohydrolase) generates glutamate from the glutamine provided by the astrocytes (Figure 1A). Glutamate synthesis occurs predominantly in neurons, although astrocytes can synthesize it through the same pathway (Hogstad et al., 1988). Neurons and astrocytes both preferentially express the mitochondrially-located, kidney-type glutaminase (GLS) isoenzyme type 1 (GLS1), although there is also the liver-type GLS2 that localizes to the mitochondrion and nucleus (Cardona et al., 2015). However, evidence suggests that neurons and astrocytes might express an isoform of glutaminase GLS1, which has not been characterized to date (Kvamme et al., 2001; Cardona et al., 2015).

Glutamate is a component of energy metabolism, which requires *de novo* synthesis to avoid an imbalance in glutamate concentrations. *De novo* glutamate synthesis occurs by the pyruvate carboxylase, which is located exclusively in astrocytic mitochondria; this enzyme metabolizes pyruvate into oxaloacetate, a precursor for α -ketoglutarate (Walker, 2014; Schousboe et al., 2019). Although neurons lack pyruvate carboxylase, they contribute to *de novo*

glutamate synthesis by the pyruvate carboxylation to malate through the malic enzyme, which is found in the cytosol and mitochondria (McKenna et al., 1995; Hassel, 2001; Amaral et al., 2016).

Another enzyme that contributes to glutamate production is aspartate aminotransferase (found in the cytosol or mitochondria), which synthesizes glutamate by reversibly transferring the α -amino group from aspartate to 2-oxoglutarate, resulting in glutamate and oxaloacetate; this enzyme uses pyridoxal 5'-phosphate as a co-factor (Figure 1B; McKenna et al., 2006; Schousboe, 2017). Neurons and astrocytes express aspartate aminotransferase, and the enzyme appears to have the same function and activity in both cell types (McKenna et al., 2006).

Astrocytes and neurons contain alanine aminotransferase in the cytoplasm and mitochondria (Ruscak et al., 1982; Waagepetersen et al., 2000), which catalyzes the reversible interconversion of alanine and α -ketoglutarate into pyruvate and glutamate (Figure 1C). Low activity of this enzyme in neurons (Westergaard et al., 1993; Erecinska et al., 1994) suggests that, within this pathway, astrocytes exert a primary control (Schousboe et al., 2013).


Ammonia concentration in the brain is regulated by the mitochondrial glutamate dehydrogenase, which catalyzes the reversible conversion of glutamate to α -ketoglutarate and ammonia, using NADH or NADPH as a co-factor (Islam et al., 2010; Plaitakis et al., 2017). Glutamate dehydrogenase expression in astrocytes varies spatially by brain region, cellularly by astrocyte type, and temporally with the developmental stage (Figure 1D; Osterberg and Wattenberg, 1962). For example, astrocytes increase glutamate dehydrogenase expression during rat hippocampus maturation (Kugler and Schleyer, 2004). Interestingly, along with increased glutamate dehydrogenase activity, astrocytes also increase the expression of GLT-1, suggesting a deeper, interconnected regulatory system of glutamatergic dynamics (Kugler and Schleyer, 2004).

Glutamate packaging

In neurons, glutamate is packaged and stored in synaptic vesicles through specific vesicular glutamate transporters (VGLUT). Currently, three subtypes of VGLUTs (VGLUT1, 2, and 3) have been described, and their distribution differs among different brain structures. VGLUT1 is present in the cerebral cortex, cerebellum, hippocampus, and thalamus (Fujiyama et al., 2001; Herzog et al., 2004). VGLUT2 is expressed in the cortex, thalamus, diencephalon, and rhombencephalon (Fremeau et al., 2001; Herzog et al., 2004). VGLUT3 is less predominant than the other two transporters and is located in the striatum, neocortex, and hippocampus (Fremeau et al., 2002).

VGLUT function depends on the electrochemical proton gradient generated across the membrane by the activity of the vacuolar H^+ -ATPase (Wolosker et al., 1996). This H^+ -ATPase activity increases the H^+ concentration inside the vesicle leading to an acidic pH. The rate of glutamate transport by VGLUT correlates inversely with the concentration of chloride ions (Cl^-) such that a low extravesicular Cl^- concentration generates high glutamate uptake, whereas high Cl^- concentration leads to gradual inhibition of glutamate uptake (Wolosker et al., 1996; Juge et al., 2006).

There appears to be a proportional relationship between VGLUT levels and glutamatergic synapse response. A study of VGLUT1-knockout mice demonstrated that the knockout reduced the



amplitude of miniature excitatory postsynaptic currents (mEPSCs), suggesting a smaller quantal size (Wojcik et al., 2004). In another study, vesicles containing a lower number of VGLUT1 showed a reduced release probability (Herman et al., 2014). Similar results have been described also in VGLUT3-knockout mice (Fasano et al., 2017).


The evidence for astrocyte expression of VGLUT is contradictory and requires further investigation to clarify the situation (for a review see Hamilton and Attwell, 2010). For example, Li et al. (2013a) describe the absence of VGLUT in mouse cortical, hippocampal, and cerebellar astrocytes. On the contrary, Ormel et al. (2012) identified VGLUT1 in astrocytic processes in the rat hippocampus, frontal cortex, and striatum. Likewise, astrocytes from postnatal rat brains express VGLUT1 and VGLUT2 (Montana et al., 2004), as well as cortical cultured astrocytes (Anlauf and Derouiche, 2005). VGLUT3 has been detected in astrocytes end-feet in microcultures of rat ventral tegmental area, substantia nigra pars compacta, and raphe nuclei (Fremeau et al., 2002). Despite the controversy about astrocytic VGLUTs, evidence indicates that astrocytes contain vesicular compartments and the molecular machinery to release glutamate in vesicular packages; this phenomenon—now known as gliotransmission—was later confirmed and contributes to neuronal information processing (see below; Bezzi et al., 2004).

Glutamate release

Once glutamate is packaged into the synaptic vesicles and stored in the synaptic bouton, it is ready to be released upon the arrival of an action potential, which will induce the opening of voltage-dependent calcium channels, increasing intracellular calcium

(Ca²⁺) concentration. The Ca²⁺ influx facilitates vesicle fusion with the plasma membrane, which releases the neurotransmitters into the synaptic cleft (Figure 2; de Wit et al., 2009). The soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) family of proteins are key components for this process. These proteins are divided into two groups, the vesicular v-SNARE, highly abundant in the vesicles, and the target t-SNARE, highly expressed in the target zone in the cellular membrane (Han et al., 2017). A trans-SNARE complex -interaction between v and t SNAREs- must be formed to release the neurotransmitter; the main v-SNAREs are synaptobrevin, synaptotagmin, syntaxin, and the main t-SNARE is SNAP-25.

SNARE expression is not limited to neurons; SNAREs support gliotransmitter release in astrocytes (Crippa et al., 2006). Cultured astrocytes express synaptobrevin II and release glutamate that is reduced by the inhibitors of the neuronal exocytosis botulinum toxin-A and botulinum toxin-C, suggesting the expression of SNAP-25 and syntaxin in astrocytes (Jefitinija et al., 1997). The vesicular exocytosis process is similar between neurons and astrocytes (Crippa et al., 2006), with the principal difference arising in the initiation of exocytosis. Neuronal exocytosis is initiated by the arrival of an action potential at the synapse, which results in membrane depolarization in addition to an influx of Ca²⁺ through transmembrane channels involved in the action potential response; the Ca²⁺ wave initiates a cascade of signaling that results in vesicular release. Astrocytes cannot generate action potentials, but the astrocytic vesicular release also requires a transient increase in intracellular Ca²⁺ concentration. Astrocytes have multiple mechanisms to accomplish this Ca²⁺ increase, including activation of ionotropic or metabotropic receptors and the subsequent inositol



1,4,5-trisphosphate (IP3) signaling cascade, activation of transient receptor potential (TRP) channels, or the release of Ca^{2+} by mitochondria (Guerra-Gomes et al., 2018); or selective activation of either purinergic receptors P2Y1 or protease-activated receptor 1 (PAR-1) that in astrocytes leads to increased intracellular Ca^{2+} concentration (Shigetomi et al., 2008).

Furthermore, recent evidence indicates that astrocytes can release glutamate by the interaction of metabotropic receptors PAR-1 with either the two-pore domain potassium channel (TREK-1) or the Bestrophin-1 (Best-1), a Ca^{2+} -activated chloride channel (Figure 2). Glutamate efflux from the intracellular space occurs when the TREK-1 intracellular domain interacts with PAR-1 allowing fast glutamate transient currents, whereas the interaction of PAR-1 and

Best-1 leads to slow transient currents (Woo et al., 2012; Lalo et al., 2021). Taken together, the evidence supports the view that both neurons and astrocytes contribute to glutamatergic signaling.

Glutamate receptors

Glutamatergic receptors are widely distributed in the different regions of the central nervous system. Neurons and astrocytes express glutamate receptors, which split into two families. Ionotropic glutamate receptors (iGluRs; Kukley et al., 2001) and metabotropic glutamate receptors (mGluRs; Schools and Kimelberg, 1999; Fiacco and McCarthy, 2004; Perea and Araque, 2007; Cavaccini et al., 2020).

iGluRs

The iGluRs family, consisting of NMDA (N-methyl-D-aspartate) receptors, and the AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazol-propionate) and KA (kainic acid) receptors, which are responsible for the excitatory transmission in the central nervous system of vertebrates. These receptors are ligand-gated ion channels that allow the movement of cations such as Na^+ , K^+ , and Ca^{2+} across the cell membrane. Structurally, iGluRs are transmembrane proteins composed of four subunits that form a central ionic pore comprised of an extracellular amino-terminal domain, an extracellular ligand-binding domain, four transmembrane domains, and an intracellular carboxyl-terminal domain (Traynelis et al., 2010).

NMDA receptor

NMDA receptors are highly expressed in the brain, and their role in physiological and pathological mechanisms have been studied extensively (Lakhan et al., 2013; Zhou et al., 2013; Intson et al., 2022). The subunits that constitute these receptors are GluN1, GluN2 [A, B, C, D], GluN3A, and GluN3B. A functional receptor contains a tetrameric assembly with two possible configurations. All receptors must have two GluN1 subunits. The remaining two subunits can be a pair of GluN2 subunits, or one GluN2 subunit together with one GluN3 subunit (Schorge et al., 2005; Ulbrich and Isacoff, 2008; Traynelis et al., 2010; Hansen et al., 2021). Moreover, mRNA editing and alternative splicing of the different NMDA subunit genes confer more complex properties to the NMDA receptors' conformation, which have been extensively reviewed by Hansen and colleagues (Hansen et al., 2021).

Full activation of NMDA receptors in neurons requires membrane depolarization to displace the Mg^{2+} ion that blocks the ion channel with the simultaneous binding of glutamate and the co-agonist, glycine. When both conditions are met, NMDA receptor activation allows Na^+ and Ca^{2+} ion influx. In addition to glycine, D-serine is an NMDA receptor co-agonist, which is supplied by astrocytes (Henneberger et al., 2010). However, recent evidence indicated that neurons also contribute to the *de novo* synthesis of D-serine (Neame et al., 2019). D-serine metabolism depends on 3-phosphoglycerate dehydrogenase (Phgdh) enzyme activity and glycine concentration (Shibasaki et al., 2017; Neame et al., 2019). Since astrocytes require Phgdh to synthesize L-serine from glucose, astrocyte *de novo* synthesis of L-serine may be a previously unrecognized regulatory mechanism for the NMDA receptors co-agonism by D-serine (Masuoka et al., 2019; Neame et al., 2019).

On the other hand, astrocytes also express NMDA receptors, as evidenced by the presence of GluN1 and GluN2 subunit mRNA (Jimenez-Blasco et al., 2015). In astrocytes, the NMDA receptors are insensitive or weakly sensitive to the blockade of Mg^{2+} ion, and their activation might even occur at negative resting potential (Verkhatsky and Kirchoff, 2007). Interestingly, the expression pattern of NMDA subunits in astrocytes varies depending on the brain region. For example, cortical astrocytes express GluN1, GluN2A/B, GluN2C, and GluN2D subunits (Conti et al., 1996; Palygin et al., 2011). GluN2B is more abundant in the Bergmann glia cells (Luque and Richards, 1995), whereas the GluN2C is predominantly expressed in the telencephalon glial cells (Alsaad et al., 2019). In hippocampal cultured astrocytes, GluN1 and GluN2 subunits are predominant

(Araque et al., 1998a). *In vivo* evaluation of the GluN2C NMDA subunit localization showed that parvalbumin-positive neurons in the globus pallidus, ventral pallidum, and substantia nigra express this subunit, whereas GluN2C in the cortex, striatum, hippocampus, and amygdala colocalizes with astrocytes' markers (Ravikrishnan et al., 2018). The variety of NMDA subunit compositions confers different activation properties to the receptors, which, in combination with the regional expression patterns, results in differences in the contribution of NMDA receptors to neuronal and astrocyte activity (Palygin et al., 2011).

AMPA receptor

An AMPA receptor is composed of four subunits from the proteins GluA1, GluA2, GluA3, and GluA4 (Traynelis et al., 2010). AMPA receptors can either be homomers or heteromers. These subunit types are differentially expressed in the nucleus accumbens, dorsal striatum, prefrontal cortex, and hippocampus (Reimers et al., 2011). Activation of AMPA receptors allows Na^+ and K^+ influx, but if the receptor conformation lacks a GluA2 subunit or contains a post-transcriptionally modified GluA2 by RNA editing at the Q/R site, the channel will also be Ca^{2+} permeable (Traynelis et al., 2010; Hansen et al., 2021). AMPA subunits interact with the transmembrane AMPA receptor regulatory protein (TARP), which modulates channel opening (Hansen et al., 2021). Activation of AMPA receptors leads to membrane depolarization, leading to the displacement of the Mg^{2+} ion that blocks the NMDA ion channel allowing its activation.

Some studies suggest the presence of AMPA receptors in glial cells (Müller et al., 1992). These receptors have been described in cortical cultured astrocytes (David et al., 1996), and a subpopulation of hippocampal astrocytes express GluA1, GluA2, GluA3, and GluA4 subunits (Matthias et al., 2003). Regarding function, it has been shown that the AMPA receptors modulate the inward-rectifier potassium channels (also known as Kir) in hippocampal astrocytes, which induces gliotransmitter release and AMPA activation in neurons (Schröder et al., 2002; Fiacco and McCarthy, 2004). Of note, AMPA receptors containing at least one GluA3 or GluA4 subunit are permeable to Ca^{2+} , as described in hippocampal astrocytes (Seifert and Steinhauser, 1995), providing a direct mechanism to raise intracellular Ca^{2+} concentrations.

KA receptor

Among the iGluRs family proteins, KA receptors comprise the least studied class (Meyerson et al., 2016). The subunits that form the KA receptors are GluK1, GluK2, GluK3, GluK4, and GluK5. The KA channel is permeable to Na^+ and K^+ . Some of the most important functions of these receptors are the regulation of synaptic activity (Fernandes et al., 2009) and neuronal plasticity (Lauri et al., 2006).

The expression of receptor subunits does indeed vary across species and brain regions. For example, the primate neocortex primarily expresses the GluK1-2-3 subunits (Huntley et al., 1993), whereas the rodent cortex expresses more GluK2 and GluK4 subunits (Herb et al., 1992). Hippocampal interneurons express KA receptors (Liu et al., 2004). Astrocytes in the hypothalamic arcuate nucleus express GluK1-3 subunits (Diano et al., 1998).

Hippocampal astrocytes express GluK2 (Matschke et al., 2015). Some KA subunits allow Ca^{2+} efflux, which in astrocytes can contribute to glutamate vesicular release or activate other Ca^{2+} -dependent signaling pathways (see Guerra-Gomes et al., 2018). Interestingly, after chemoconvulsive status epilepticus of temporal lobe epilepsy, CA1 hippocampal reactive astrocytes expressed GluK1, GluK2/3, GluK4, and GluK5 (Vargas et al., 2013). Of these, GluK1 and GluK5 expression in astrocytes persist during the presence of spontaneous seizures, suggesting that KA receptors in astrocytes might contribute to the pathophysiology of epilepsy (Vargas et al., 2013).

In general, iGluRs are a key component for synaptic activity and neuronal processing, however, more research is required to elucidate the contribution of each iGluRs in astrocytes and how it contributes to glutamatergic neurotransmission.

mGluRs

mGluRs are coupled to G-proteins and modulate slow synaptic transmission through second messengers. To date, eight mGluRs (mGluR1–8) have been described and divided into three groups designated I, II, and III according to similarities in their distinctive features: gene sequence, pharmacological properties, and intracellular signaling mechanisms (Sladeczek et al., 1985).

The group I receptors, mGluR1 and mGluR5, are associated with intracellular Ca^{2+} signaling, phospholipase C, and these receptors are mainly activated by 3,5-dihydroxyphenylglycine (DHPG). Group II includes mGluR2 and mGluR3, which are negatively coupled to adenylate cyclase and are selectively activated by LY379268. Finally, group III contains mGluR4, mGluR6, mGluR7, and mGluR8 receptors, which, like group II, are negatively coupled to the adenylate cyclase (Sugiyama et al., 1987; Masu et al., 1991).

Receptors from the Group I mGluRs are more widespread in the brain. They are expressed in neurons from the olfactory bulb, cerebral cortex, globus pallidus, lateral septum, cerebellar Purkinje cells, and thalamic nuclei (Crupi et al., 2019). Group II mGluRs are expressed in the olfactory bulb and cerebellar cortex (for more details see Crupi et al., 2019). In contrast, astrocytes showed a predominance of mGluR1, mGluR3, and mGluR5 receptors, which have been described thus far in the hippocampus and cerebral cortex (Schools and Kimelberg, 1999; Sun et al., 2013; Spampinato et al., 2018). Using electron microscopy and immunohistochemistry, mGluR2 and mGluR3 have been identified in astrocytes in the rat ventrobasal thalamus (Mineff and Valtschanoff, 1999).

The activation of mGluR3 and mGluR5 increases Ca^{2+} intracellular concentration, triggering vesicular glutamate release in neurons and astrocytes, which influences synaptic activity and plasticity (Fiacco and McCarthy, 2004; Perea and Araque, 2007; Cavaccini et al., 2020). Specific activation of metabotropic group II receptors in astrocyte cultures increases the expression of GLAST (Gegelashvili et al., 2000). Likewise, activating the mGluR5 receptor increases glutamate uptake through increased expression of GLT-1 and GLAST transporters (Vermeiren et al., 2005). However, contradicting results have been observed during the activation of group I mGluRs, which reduces the expression of GLAST (Gegelashvili et al., 2000). Therefore, more studies are needed to understand the specific role of each mGluR in astrocytic glutamate transporter expression and function.

Glutamate transporters

Glutamate transporters, also known as excitatory amino acid transporters (EAATs), maintain optimal extracellular glutamate concentration; these transporters belong to the solute carrier (SLC) family 1 (high-affinity glutamate transporters; He et al., 2009). These proteins are expressed in neurons and glial cells, especially astrocytes, and are responsible for the bulk of glutamate uptake (Rothstein et al., 1996) by the co-transport of glutamate, Na^+ (three molecules), H^+ (one molecule), and counter-transport of K^+ (one molecule; Levy et al., 1998).

Five EAATs have been identified in humans. In rodents, these transporters were named excitatory amino acid carrier 1 EAAC1/EAAT3, GLAST/EAAT1, GLT-1/EAAT2, excitatory amino acid transporter 4 (EAAT4), and excitatory amino acid transporter 5 (EAAT5; Figure 2; Kanai and Hediger, 1992; Pines et al., 1992; Storck et al., 1992; Fairman et al., 1995; Rothstein et al., 1996; Arriza et al., 1997).

Neurons in the rat cerebral cortex, hippocampus, cerebellum, and spinal cord express the EAAC1 transporters (Kanai et al., 1995; Shashidharan et al., 1997). Interestingly, this transporter is mainly involved in anion conductance and uptake of cysteine, a precursor of glutathione synthesis (Lee et al., 2020). EAAT4 is highly expressed in cerebellar Purkinje cells (Magi et al., 2019). However, it is also found in the fore- and mid-brain and the somatosensory cortex (Massie et al., 2008; de Vivo et al., 2010). EAAT5 is mainly expressed in the retina (Arriza et al., 1997).

GLAST and GLT-1 are highly expressed in astrocytes of the hippocampus, striatum, and cerebral cortex and oversee glutamate uptake at the synapse (Levy et al., 1993; Lehre et al., 1995; Bergles and Jahr, 1997; Mennerick et al., 1998). According to studies focused on evaluating the subcellular distribution of GLAST and GLT-1, both transporters are highly expressed in hippocampal astrocytes with a predominant presence of GLT-1 in the filopodium and perivascular end-feet, and GLAST is mostly present in the soma and processes (Schreiner et al., 2014; Radulescu et al., 2022). However, the GLT-1 isoforms (GLT-1a and b) are expressed in neurons from the hippocampus, cerebral cortex, striatum, thalamus, and midbrain (Chen et al., 2002, 2004; Berger et al., 2005). Astrocyte processes express more GLT-1a mRNA, whereas GLT-1b mRNA has been detected mainly in the cell body (Berger et al., 2005). In neurons, GLT-1a protein expression in axons, spines, and dendrites contributes to glutamate reuptake in the excitatory terminals (Chen et al., 2004). It has been suggested that GLT-1 in neurons provides glutamate as a substrate for energy metabolism and mitochondrial functionality (Petr et al., 2015; McNair et al., 2019).

Astrocytes are the main regulators of extracellular glutamate concentration through the GLT-1 and GLAST glutamate transporters; expression of these transporters is regulated by neuronal activity (Swanson et al., 1997; Perego et al., 2000). Interestingly, besides neurons, brain endothelial cells can also induce GLT-1 expression through Notch signaling (Lee et al., 2017).

In addition to EAATs, the SLC7A11/xCT transporter is a cystine/glutamate antiporter, which transports a cystine into the cell while exchanging for glutamate (1–1 ratio), in a sodium-independent fashion; therefore, contributing to astrocyte glutamate release (Bannai, 1986). It consists of two subunits, the light subunit (SLC7A11) and the heavy subunit (SLC3A2). Whereas the light subunit is responsible for the active transport of cystine and glutamate, the heavy subunit is necessary for intracellular trafficking

and proper membrane arrangement of the transporter (Nakamura et al., 1999; Shin et al., 2017). The SLC7A11/xCT transporter takes up cystine and, inside the cell, cystine will be converted to cysteine, the main precursor for the antioxidant glutathione (Conrad and Sato, 2012). SLC7A11/xCT is highly expressed in the human brain (Sato et al., 1999). In mice, SLC7A11/xCT is prominently expressed in the hippocampus, cortex, hypothalamus, and dentate gyrus (Sato et al., 2002). The SLC7A11/xCT expression occurs mainly in glial cells (Re et al., 2006), including astrocytes (Ottestad-Hansen et al., 2018). SLC7A11/xCT is essential to avoid oxidative damage (Lewerenz et al., 2012), probably due to its link with glutathione synthesis. It follows that blocking SLC7A11/xCT leads to an increase in oxidative stress and astrocyte death (Chen et al., 2000), a process known as oxidative glutamate toxicity (Schubert and Piasecki, 2001).

Interestingly, *Drosophila* xCT gene-knockout reduced the extracellular ambient glutamate concentration by 50%, suggesting that the xCT transporter is essential for extracellular glutamate regulation (Augustin et al., 2007). Also, the use of sulfasalazine, an xCT inhibitor, reduces the NMDA-induced current by 66.8% in mouse hippocampus slices, indicating that glutamate release through xCT contributes to neuronal activation (Koh et al., 2022). In addition, xCT deletion in mice induces an age-dependent anxiety-like behavior (Bentea et al., 2015). In a related experiment, exposing the astrocytoma-derived cell line (1321N1) to peroxide increased both ambient glutamate concentrations and the population of xCT transporters, suggesting that xCT activity contributes to glutamate release and accumulation (Kazama et al., 2020). Further experiments are necessary to clarify whether the release of glutamate by the xCT transporter contributes to the activation of iGluRs and mGluRs in neurons or astrocytes and if it contributes to pathological processes *in vivo*.

Role of astrocytes in glutamatergic neurotransmission during physiological conditions

Astrocyte function was initially thought to support neuronal activity or protect neurons from excitotoxicity. However, later studies suggested that astrocytes can directly or indirectly modulate synaptic neuronal activity (Figure 2; Nedergaard, 1994; Beppu et al., 2021) and influence behavior (Lyon and Allen, 2022). The first level of regulation is glutamate uptake by astrocytic transporters as they regulate the neurotransmitter levels in the synaptic cleft; these transporters indirectly modulate neuronal transmission (Jabaudon et al., 1999), neuronal activity (Estrada-Sánchez et al., 2019) and survival (Estrada-Sánchez et al., 2007, 2019). Glutamate uptake can also regulate the availability of glutamine to synthesize glutamate. As mentioned in an earlier section, once astrocytes take up glutamate, it can be metabolized into α -ketoglutarate by glutamate dehydrogenase or into glutamine through amidation of glutamate by the glutamine synthetase (as part of the glutamate/glutamine cycle; Laake et al., 1999; Islam et al., 2010). As this enzyme is highly expressed in astrocytes, they are considered the major glutamine reservoir and an important source of precursor for the metabolism of glutamate and gamma-aminobutyric acid (GABA; Hamberger et al., 1979; Norenberg and Martinez-Hernandez, 1979). Therefore, astrocytes also might regulate glutamatergic

neuronal dynamics by the amount of glutamine released into the synaptic cleft.

Astrocytes can also regulate neuronal activity by releasing gliotransmitters such as glutamate, ATP, D-serine, or GABA, also known as gliotransmission. Once the gliotransmitter is released, it activates its target receptors and, consequently, generates responses in the same astrocyte (autocrine response) or nearby cells, including neurons (Lapato and Tiwari-Woodruff, 2018; Savtchouk and Volterra, 2018; Beppu et al., 2021; Sherwood et al., 2021a). In addition to gliotransmission, astrocytes contribute to neuronal activity by regulating the availability of NMDA receptor co-agonists (glycine and D-serine; Sherwood et al., 2021a). However, more studies are needed to better understand this process's physiological and pathological implications. For a more detailed description of this topic, see Sherwood et al. (2021a). Also, astrocytes can release active molecules through hemichannels (Lee et al., 2011; Montero and Orellana, 2015; Lalo et al., 2021). This topic is beyond the scope of this review, but for more information refer to Sahlender et al. (2014), Montero and Orellana (2015), and Caudal et al. (2020).

Experiments using electrophysiology, optical imaging, and molecular biology demonstrated that astrocytes respond to neurotransmitters. Activation of mGluR initiates a cellular signaling cascade that increases intracellular Ca^{2+} concentration on a timescale of about 50–200 ms (Batchelor and Garthwaite, 1997; Marcaggi et al., 2009), in contrast to the comparatively fast ionotropic receptors that take approximately 1–10 ms in neurons to initiate the same response (Traynelis et al., 2010; Reiner and Levitz, 2018). iGluRs are fast-acting because extracellular Ca^{2+} directly enters the cell through the open channel, although a significant Ca^{2+} concentration rise requires a substantial number of simultaneously open iGluR channels. The activation of mGluR initiates a cellular signaling cascade that amplifies the input signal, albeit at the cost of response time; the activated mGluR activates the phospholipase C/IP3 pathway, which then generates the release of Ca^{2+} from the endoplasmic reticulum (Decrock et al., 2013; Rodriguez-Prados et al., 2020). This intracellular source of Ca^{2+} induces gliotransmitter release by Ca^{2+} -dependent exocytosis (Bezzi et al., 1998, 2004; Zhang et al., 2004; Mothet et al., 2005; Crippa et al., 2006; Woo et al., 2012; Li et al., 2013b; Navarrete et al., 2013; Heller et al., 2020; Takata-Tsuji et al., 2021). Gliotransmitter release in turn affects neuronal functioning, forming a feedback loop.

Although astrocytes express both iGluRs and mGluRs, intracellular Ca^{2+} concentration rises mainly due to the activation of mGluRs rather than iGluRs (Conti et al., 1996; Schools and Kimelberg, 1999). However, we cannot dismiss the contribution of iGluRs in astrocytes for two reasons. First, the temporal dynamics of iGluRs are approximately an order of magnitude faster than mGluRs. Second, the presence of certain subunits confers different Ca^{2+} permeability to iGluRs (Seifert et al., 1997; Brand-Schieber and Werner, 2003; Brand-Schieber et al., 2004; Palygin et al., 2010). The subunits that increase Ca^{2+} permeability are GluN3 subunits for NMDA (Cull-Candy et al., 2001; Kvist et al., 2013), GluA2 for AMPA (Traynelis et al., 2010), and GluK3-4 for KA (Burnashev et al., 1995). The ratio of iGluRs with and without increased Ca^{2+} permeability in astrocytes may constitute a regulatory mechanism to modulate Ca^{2+} influx into the cell, affecting Ca^{2+} -dependent pathways.

The release of glutamate from astrocytes contributes to NMDA-related long-term depression. This mechanism is initiated

by cannabinoid receptor type 1 (CB1) activation of astrocytes, which increases Ca^{2+} concentration and produces astrocytic glutamate release. Increased extracellular glutamate activates NMDA receptors, promoting internalization of AMPA receptors (Han et al., 2012; Min and Nevian, 2012) in the cortex and hippocampus. In the cortex, the activation of astrocytic CB1 receptors by exogenous cannabinoids impairs spatial working memory (Han et al., 2012; Min and Nevian, 2012).

In the striatum, two subpopulations of astrocytes may selectively regulate the response for dopamine D1 vs. D2 medium spiny neuron (MSN) subpopulations. Astrocytic CB1 activation elicits astrocytic glutamate release that specifically induces activity from either D1 or D2 MSNs, but not both (Martín et al., 2015). This finding suggests nuanced organization and interaction of neurons and astrocytes in the striatum.

On the other hand, Beppu et al. (2021) demonstrated that Bergmann glial cells amplify excitatory neuronal signals in the cerebellar cortex by releasing glutamate through a mechanism involving bicarbonate efflux and resultant intracellular acidification, a mechanism sensitive to the inhibition of volume-regulated ion channels. Although this is a novel and exciting mechanism by which astrocytes can regulate the neuronal activity, more experiments are needed to elucidate its specific molecular components and their contribution during physiological conditions *in vivo*. Also, this opens the question of whether this is a mechanism restricted to the cerebellar cortex or if it also occurs in other brain areas.

Interneurons also play an important role in astrocyte modulation of neuronal activity. Interneurons are locally projecting neurons that regulate neuronal activity levels through inhibitory signaling that counteracts excitatory (e.g., glutamatergic) signaling, and helps to prevent runaway excitatory cascades. For example, in the hippocampus, stimulation of inhibitory GABAergic interneurons activates GABA_B receptors in astrocytes, which subsequently triggers increased Ca^{2+} waves in the surrounding astrocytes, potentiating pyramidal inhibition. This effect is blocked by glutamatergic antagonist CNQX [cyanquinoxaline (6-cyano-7-nitroquinoxaline-2,3-dione)] and AP5 (2-amino-5-phosphopentanoic acid), suggesting that interneuron-astrocyte-mediated potentiated inhibition of pyramidal neurons depends on astrocyte-mediated glutamate release (Kang et al., 1998). Also, in the hippocampus, Perea et al. (2016) described that, besides activating GABA_B receptors in astrocytes, GABA release by interneurons also activates GABA_A receptors in presynaptic neurons, which inhibited synaptic activity. However, when the interneuron leads to astrocyte-mediated glutamate release, presynaptic activation of mGluR 1/5 receptors contributes to synaptic potentiation (Kang et al., 1998; Perea et al., 2016). In the mouse cerebellar cortex, activation of purinergic P2Y1 receptors and AMPA receptors in Bergmann glial cells leads to glutamate-vesicular release that activates NMDA receptors in the interneurons, enhancing the inhibitory synaptic input to Purkinje cells (Rudolph et al., 2016).

Shen et al. (2017) described that autocrine activation of P2Y1 purinergic receptors in astrocytes modulates the release of glutamate mediated by the Ca^{2+} -dependent chloride channel Best-1 and the subsequent activation of extra-synaptic NMDA receptors in neurons. Related studies evaluated the effect of astrocytic Ca^{2+} -dependent glutamate release on the activity of neuronal extra-synaptic NMDA receptors (Le Meur et al., 2007; Shen et al., 2017; Koh

et al., 2021). These receptors contain the GluN2B subunit and their activation produce a slow inward current with an amplitude of 18–477 pA, with a rise time of 13–332 ms, and decay times of 72–1,630 ms. The presence of extra-synaptic NMDA currents directly depends on astrocytes' intracellular Ca^{2+} concentrations; they decrease when Ca^{2+} signaling is abolished and increase when intracellular Ca^{2+} concentration rises (Araque et al., 1998b; Pirttimäki et al., 2011; Perea et al., 2014); these neuronal currents are generated by astrocyte activity in the hippocampus, cortex (Gomez-Gonzalo et al., 2017, 2018), and nucleus accumbens (Corkrum et al., 2019).

Moreover, Santello et al. (2011) showed that the P2Y1 activation and consequent Ca^{2+} -dependent glutamate release in astrocytes from the dentate gyrus also involve the tumor necrosis factor- α (TNF α), which at physiological concentrations (pM range), favors the adequate exocytosis of astrocytic glutamatergic vesicles. Interestingly, glutamate released activates presynaptic NMDA receptors, particularly expressing the GluN3A subunit with a low voltage-dependent Mg^{2+} block (Savtchouk et al., 2019). In an excellent review, Di Castro and Volterra (2022) describe how this mechanism might be relevant within the entorhinal cortex-dentate gyrus circuit involved in memory processing.

Additional evidence beyond the hippocampus indicates that astrocytes have a functional role in behavior regulation. For example, astrocytes in the suprachiasmatic nucleus increase Ca^{2+} signaling at night. This signaling is highly related to circadian behavior regulated by glutamate release (Brancaccio et al., 2019). Blum et al. (2021) demonstrated in *Drosophila melanogaster* through *in vivo* two-photon experiments that increased astrocyte Ca^{2+} activity correlates with sleep needs. Another behavior modulated by astrocytes is feeding (Sweeney et al., 2016; Varela et al., 2021). Specific stimulation of astrocytes from mice's medial basal hypothalamus suppresses food intake (Sweeney et al., 2016). Finally, mice lacking astrocyte glucocorticoid receptors in the amygdala show attenuated anxiety behaviors in the open field behavioral test and fear memory (Wiktorowska et al., 2021), demonstrating a direct involvement of astrocytes in fear memory and anxiety.

The extensive evidence reflects the key role of Ca^{2+} -mediated signaling in astrocytes, which requires stimulation of IP₃ receptors (IP₃R). There are three subtypes of IP₃R in mammals (IP₃R1, IP₃R2, and IP₃R3) and among them, IP₃R2 was widely accepted as the only functional subtype in astrocytes (Sherwood et al., 2021b). However, recent evidence showed that IP₃R1 and IP₃R3 are also present in astrocytes and participate in Ca^{2+} -mediated signaling, especially in astrocytic processes (Sherwood et al., 2017, 2021b). This new finding raises questions about the functional role and subcellular localization of the different IP₃R subtypes in astrocytes during gliotransmission and its relevance during physiological processes.

As a whole, this evidence indicates that gliotransmission modulates the activity of astrocytes and neuronal circuits (projecting and interneurons) and appears to be a widespread mechanism in the brain. Besides glutamate, ATP purinergic receptors emerge as a key component that triggers astrocytes' modulation of neuronal circuits. Although the current evidence indicates that gliotransmission can influence behavior, more research is needed to dissect the different roles of astrocytes in shaping animal behavior.

Role of astrocytes in glutamatergic neurotransmission during pathological conditions

Alterations in the glutamate transporters are related to neurodegenerative diseases (Pajarillo et al., 2019). *Postmortem* brains of Alzheimer's disease patients showed a reduction in EAAT1 and EAAT2 (Masliah et al., 2000). Epilepsy is associated with decreased EAAT2 (Tanaka et al., 1997). In the intrahippocampal kainic acid model of temporal lobe epilepsy, GLT-1 and GLAST expression increase early after the treatment, suggesting that dysregulation in the expression of astrocyte glutamate transporters could contribute to the development of epilepsy. However, the accuracy of this hypothesis has not yet been determined (Peterson and Binder, 2019). In Huntington's disease, an inherited neurodegenerative disorder, reduced expression of EAAT2 was observed in *postmortem* brain samples (Faideau et al., 2010). In the R6/2 Huntington's disease transgenic mouse model, decreased content of GLT-1 and GLAST correlates with increased vulnerability to glutamate-induced toxicity (Estrada-Sánchez et al., 2009, 2010). Likewise, cortical pyramidal neurons in the R6/2 mice are more vulnerable to glutamate-mediated paroxysmal activity during the inhibition of both GLT-1 and GLAST transporters (Estrada-Sánchez et al., 2019).

To date, exploration of how gliotransmission contributes to neural information processing and behavior focused mostly on physiological conditions. Less is known about how changes in gliotransmission contribute to the pathology of neurological, neuropsychiatric, or neurodegenerative conditions. However, data from physiological studies point out that astrocytes in the striatum can modulate differentially D1 or D2 MSNs (see above), and its dysfunction may contribute to diseases like Parkinson's and Huntington's disease. D1 and D2 MSNs comprise key components of the brain circuits that control movement and, at the behavioral level, both diseases involve substantial alterations in movement control. At the cellular level, D1/D2 MSN impairment contributes to neuropathology, and a distinct line of evidence demonstrates altered astrocyte functioning (for a review, see Estrada-Sánchez et al., 2017). The results from Martín et al. (2015) suggest that these two lines of evidence from disease pathology are connected to healthy tissue function (Martín et al., 2015); further understanding of how this system works may yield promising pathways for future therapeutic interventions for these diseases.

Another possible contribution of gliotransmission to neuropathology might be through the activation of extra-synaptic NMDA receptors since the NMDA-mediated response in neurons depends on its subcellular localization. Whereas activation of synaptic NMDA receptors leads to survival pathways, activation of extra-synaptic NMDA receptors initiates neuronal death cascades (Kaufman et al., 2012). In fact, it has been documented that activation of extra-synaptic NMDA receptors contributes to the neurodegenerative processes described in Huntington's disease and ischemia (Hardingham and Bading, 2010; Milnerwood et al., 2012). Because activation of extra-synaptic NMDA receptors indicates astrocytic Ca^{2+} -dependent glutamate release, it is important to evaluate the possible role of astrocytes in the balance/imbalance between

synaptic and extra-synaptic glutamatergic receptors activation and whether this contributes to survival or neuronal death pathways.

The description of three functional IP₃R subtypes in astrocytes raises new questions about their role in pathological states. Recently, it was shown that the IP₃R1 subtype has a key role in chronic itching (Shiratori-Hayashi et al., 2021) and the absence of IP₃R2 in astrocytes generates autism spectrum disorder-like behaviors (Wang et al., 2021).

More studies are needed to fill the knowledge gaps about the contribution of gliotransmission during the pathological process of neurodegenerative, psychiatric, and neuropathological conditions.

Perspectives

The evidence reviewed strongly suggests a complex functional interaction between neurons and astrocytes. The extent to which astrocytes modulate the synapse could vary depending on the brain region, influencing its information processing and behavioral output. To date, most of the information on gliotransmission has been centered in the hippocampus, and less is known about gliotransmission in other areas of the brain. Similarly, most studies have focused on physiological conditions, and although there is evidence that astrocytes contribute to the neuropathological process, the precise role of astrocytes during neuropathological processing is still to be determined. The review also suggests extensive opportunities for further research, including the specific contribution of each gliotransmitter described to date and perhaps the identification of new gliotransmitters, their synthesis, and release systems. It is also important to better understand the effects of gliotransmitters on neighboring cells, including the same astrocyte or afferent, efferent neurons, interneurons, and microglia cells. Finally, additional studies will clarify the functional interconnection among different signaling pathways in a tripartite synapse, such as glutamatergic, purinergic, and GABAergic, during physiological and pathological conditions. Current evidence is limited to *in vitro* and brain slice experiments, which limits our understanding of the functional role of all these components *in vivo*.

Conclusions

The development of new experimental tools has widened our understanding of the synapse, where astrocytes emerge as a complex contributor. The evidence indicates that astrocyte release of gliotransmitters such as glutamate, ATP, D-serine, and GABA can activate, potentiate, or inhibit the activity of projection neurons, interneurons, or other astrocytes. Furthermore, astrocytes can influence neuronal synaptic modification through effects on long-term depression and potentiation. It is likely that more than one gliotransmitter coexists within the same astrocyte, providing a new degree of complexity to astrocyte modulatory activity. These mechanisms suggest additional layers of information-processing capability, enabled by astrocytes that extend beyond the traditional focus on neurons as the information-processing cells of the brain. More studies are needed to understand how astrocytes modulate neuron-astrocyte network activity across the brain during

physiological and pathological conditions. Finally, deepening the understanding of the functional dynamics between gliotransmitter signaling in neurons and astrocytes will widen the therapeutic targets for neuropathological conditions and neurodegenerative diseases, including ischemia, stroke, and Huntington's disease.

Author contributions

AOC-S and AME-S conceived the review topic. AOC-S, VMR-R, TBM-B, AP-S, AGH, DPP-P, AMG, and AME-S prepared the first draft and figures. All authors contributed to the article and approved the submitted version.

Funding

This work received support from the Consejo Nacional de Ciencia y Tecnología (CONACYT) FOSEC SEP-INVESTIGACIÓN BÁSICA A1-S-26479, and the International Society for Neurochemistry-CATEGORY 1C: Return Home grant to AME-S. CONACYT also awarded a master's fellowship to AOC-S (753050), and doctoral fellowship to AOC-S (800824), TBM-B (714872), and a postdoctoral fellowship to VMR-R (626561). AMG was funded by PR-INBRE (NIH-NIGMS P20GM103475).

References

- Allen, N. J., and Eroglu, C. (2017). Cell biology of astrocyte-synapse interactions. *Neuron* 96, 697–708. doi: 10.1016/j.neuron.2017.09.056
- Alsaad, H. A., DeKorver, N. W., Mao, Z., Dravid, S. M., Arikath, J., and Monaghan, D. T. (2019). In the telencephalon, GluN2C NMDA receptor subunit mRNA is predominately expressed in glial cells and GluN2D mRNA in interneurons. *Neurochem. Res.* 44, 61–77. doi: 10.1007/s11064-018-2526-7
- Amaral, A. I., Hadera, M. G., Tavares, J. M., Kotter, M. R., and Sonnewald, U. (2016). Characterization of glucose-related metabolic pathways in differentiated rat oligodendrocyte lineage cells. *Glia* 64, 21–34. doi: 10.1002/glia.22900
- Anlauf, E., and Derouiche, A. (2005). Astrocytic exocytosis vesicles and glutamate: a high-resolution immunofluorescence study. *Glia* 49, 96–106. doi: 10.1002/glia.20094
- Anlauf, E., and Derouiche, A. (2013). Glutamine synthetase as an astrocytic marker: its cell type and vesicle localization. *Front. Endocrinol. (Lausanne)* 4:144. doi: 10.3389/fendo.2013.00144
- Araque, A., Sanzgiri, R. P., Parpura, V., and Haydon, P. G. (1998a). Calcium elevation in astrocytes causes an NMDA receptor-dependent increase in the frequency of miniature synaptic currents in cultured hippocampal neurons. *J. Neurosci.* 18, 6822–6829. doi: 10.1523/JNEUROSCI.18-17-06822.1998
- Araque, A., Parpura, V., Sanzgiri, R. P., and Haydon, P. G. (1998b). Glutamate-dependent astrocyte modulation of synaptic transmission between cultured hippocampal neurons. *Eur. J. Neurosci.* 10, 2129–2142. doi: 10.1046/j.1460-9568.1998.00221.x
- Araque, A., Parpura, V., Sanzgiri, R. P., and Haydon, P. G. (1999). Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci.* 22, 208–215. doi: 10.1016/s0166-2236(98)01349-6
- Arriza, J. L., Eliasof, S., Kavanaugh, M. P., and Amara, S. G. (1997). Excitatory amino acid transporter 5, a retinal glutamate transporter coupled to a chloride conductance. *Proc. Natl. Acad. Sci. U S A* 94, 4155–4160. doi: 10.1073/pnas.94.8.4155
- Augustin, H., Grosjean, Y., Chen, K., Sheng, Q., and Featherstone, D. E. (2007). Nonvesicular release of glutamate by glial xCT transporters suppresses glutamate receptor clustering *in vivo*. *J. Neurosci.* 27, 111–123. doi: 10.1523/JNEUROSCI.4770-06.2007
- Bannai, S. (1986). Exchange of cystine and glutamate across plasma membrane of human fibroblasts. *J. Biol. Chem.* 261, 2256–2263. doi: 10.1016/S0021-9258(17)35926-4
- Batchelor, A. M., and Garthwaite, J. (1997). Frequency detection and temporally dispersed synaptic signal association through a metabotropic glutamate receptor pathway. *Nature* 385, 74–77. doi: 10.1038/385074a0
- Behar, T. N., Scott, C. A., Greene, C. L., Wen, X., Smith, S. V., Maric, D., et al. (1999). Glutamate acting at NMDA receptors stimulates embryonic cortical neuronal migration. *J. Neurosci.* 19, 4449–4461. doi: 10.1523/JNEUROSCI.19-11-04449.1999
- Bentea, E., Demuyser, T., Van Liefvering, J., Albertini, G., Deneyer, L., Nys, J., et al. (2015). Absence of system xc⁻ in mice decreases anxiety and depressive-like behavior without affecting sensorimotor function or spatial vision. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 59, 49–58. doi: 10.1523/JNEUROSCI.1672-22.2022
- Beppu, K., Kubo, N., and Matsui, K. (2021). Glial amplification of synaptic signals. *J. Physiol.* 599, 2085–2102. doi: 10.1113/JP280857
- Berger, U. V., DeSilva, T. M., Chen, W., and Rosenberg, P. A. (2005). Cellular and subcellular mRNA localization of glutamate transporter isoforms GLT1a and GLT1b in rat brain by *in situ* hybridization. *J. Comp. Neurol.* 492, 78–89. doi: 10.1002/cne.20737
- Bergles, D. E., and Jahr, C. E. (1997). Synaptic activation of glutamate transporters in hippocampal astrocytes. *Neuron* 19, 1297–1308. doi: 10.1016/s0896-6273(00)80420-1
- Bezzi, P., Carmignoto, G., Pasti, L., Vesce, S., Rossi, D., Rizzini, B. L., et al. (1998). Prostaglandins stimulate calcium-dependent glutamate release in astrocytes. *Nature* 391, 281–285. doi: 10.1038/34651
- Bezzi, P., Gundersen, V., Galbete, J. L., Seifert, G., Steinhäuser, C., Pilati, E., et al. (2004). Astrocytes contain a vesicular compartment that is competent for regulated exocytosis of glutamate. *Nat. Neurosci.* 7, 613–620. doi: 10.1038/nn1246
- Blum, I. D., Keleş, M. F., Baz, E. S., Han, E., Park, K., Luu, S., et al. (2021). Astroglial calcium signaling encodes sleep need in *Drosophila*. *Curr. Biol.* 31, 150–162.e7. doi: 10.1016/j.cub.2020.10.012
- Brancaccio, M., Edwards, M. D., Patton, A. P., Smyllie, N. J., Chesham, J. E., Maywood, E. S., et al. (2019). Cell-autonomous clock of astrocytes drives circadian behavior in mammals. *Science* 363, 187–192. doi: 10.1126/science.aat4104
- Brand-Schieber, E., Lowery, S. L., and Werner, P. (2004). Select ionotropic glutamate AMPA/kainate receptors are expressed at the astrocyte-vessel interface. *Brain Res.* 1007, 178–182. doi: 10.1016/j.brainres.2003.12.051
- Brand-Schieber, E., and Werner, P. (2003). AMPA/kainate receptors in mouse spinal cord cell-specific display of receptor subunits by oligodendrocytes and

Acknowledgments

We would like to thank Manuel Estrada for his help during figure preparation and the IPICYT for providing access to academic journals for the review.

Conflict of interest

AGH was employed by the company Intelligent Systems Laboratory, HRL Laboratories, LLC.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- astrocytes and at the nodes of Ranvier. *Glia* 42, 12–24. doi: 10.1002/glia.10136
- Burnashev, N., Zhou, Z., Neher, E., and Sakmann, B. (1995). Fractional calcium currents through recombinant GluR channels of the NMDA, AMPA and kainate receptor subtypes. *J. Physiol.* 485, 403–418. doi: 10.1113/jphysiol.1995.sp020738
- Cardona, C., Sanchez-Mejias, E., Davila, J. C., Martin-Rufian, M., Campos-Sandoval, J. A., Vitorica, J., et al. (2015). Expression of Glu and Gls2 glutaminase isoforms in astrocytes. *Glia* 63, 365–382. doi: 10.1002/glia.22758
- Caudal, L. C., Gobbo, D., Scheller, A., and Kirchhoff, F. (2020). The paradox of astroglial Ca²⁺ signals at the interface of excitation and inhibition. *Front. Cell. Neurosci.* 14:609947. doi: 10.3389/fncel.2020.609947
- Cavaccini, A., Durkee, C., Kofuji, P., Tonini, R., and Araque, A. (2020). Astrocyte signaling gates long-term depression at corticostriatal synapses of the direct pathway. *J. Neurosci.* 40, 5757–5768. doi: 10.1523/JNEUROSCI.2369-19.2020
- Chen, W., Aoki, C., Mahadomrongkul, V., Gruber, C. E., Wang, G. J., Blitzblau, R., et al. (2002). Expression of a variant form of the glutamate transporter GLT1 in neuronal cultures and in neurons and astrocytes in the rat brain. *J. Neurosci.* 22, 2142–2152. doi: 10.1523/JNEUROSCI.22-06-02142.2002
- Chen, C. J., Liao, S. L., and Kuo, J. S. (2000). Gliotoxic action of glutamate on cultured astrocytes. *J. Neurochem.* 75, 1557–1565. doi: 10.1046/j.1471-4159.2000.0751557.x
- Chen, W., Mahadomrongkul, V., Berger, U. V., Bassan, M., DeSilva, T., Tanaka, K., et al. (2004). The glutamate transporter GLT1a is expressed in excitatory axon terminals of mature hippocampal neurons. *J. Neurosci.* 24, 1136–1148. doi: 10.1523/JNEUROSCI.1586-03.2004
- Conrad, M., and Sato, H. (2012). The oxidative stress-inducible cystine/glutamate antiporter, system x (c) (-): cystine supplier and beyond. *Amino Acids* 42, 231–246. doi: 10.1007/s00726-011-0867-5
- Conti, F., DeBiasi, S., Minelli, A., and Melone, M. (1996). Expression of NR1 and NR2A/B subunits of the NMDA receptor in cortical astrocytes. *Glia* 17, 254–258. doi: 10.1002/(SICI)1098-1136(199607)17:3<254::AID-GLIA7>3.0.CO;2-0
- Corkrum, M., Rothwell, P. E., Thomas, M. J., Kofuji, P., and Araque, A. (2019). Opioid-mediated astrocyte-neuron signaling in the nucleus accumbens. *Cells* 8:586. doi: 10.3390/cells8060586
- Crippa, D., Schenk, U., Francolini, M., Rosa, P., Verderio, C., Zonta, M., et al. (2006). Synaptobrevin2-expressing vesicles in rat astrocytes: insights into molecular characterization, dynamics and exocytosis. *J. Physiol.* 570, 567–582. doi: 10.1113/jphysiol.2005.094052
- Crupi, R., Impellizzeri, D., and Cuzzocrea, S. (2019). Role of metabotropic glutamate receptors in neurological disorders. *Front. Mol. Neurosci.* 12:20. doi: 10.3389/fnmol.2019.00020
- Cull-Candy, S., Brickley, S., and Farrant, M. (2001). NMDA receptor subunits: diversity, development and disease. *Curr. Opin. Neurobiol.* 11, 327–335. doi: 10.1016/s0959-4388(00)00215-4
- David, J. C., Yamada, K. A., Bagwe, M. R., and Goldberg, M. P. (1996). AMPA receptor activation is rapidly toxic to cortical astrocytes when desensitization is blocked. *J. Neurosci.* 16, 200–209. doi: 10.1523/JNEUROSCI.16-01-00200.1996
- de Vivo, L., Melone, M., Bucci, G., Rothstein, J. D., and Conti, F. (2010). Quantitative analysis of EAAT4 promoter activity in neurons and astrocytes of mouse somatic sensory cortex. *Neurosci. Lett.* 474, 42–45. doi: 10.1016/j.neulet.2010.03.003
- de Wit, H., Walter, A. M., Milosevic, I., Gulyas-Kovacs, A., Riedel, D., Sorensen, J. B., et al. (2009). Synaptotagmin-1 docks secretory vesicles to syntaxin-1/SNAP-25 acceptor complexes. *Cell* 138, 935–946. doi: 10.1016/j.cell.2009.07.027
- Decrock, E., De Bock, M., Wang, N., Gadicherla, A. K., Bol, M., Delvaeye, T., et al. (2013). IP3, a small molecule with a powerful message. *Biochim. Biophys. Acta* 1833, 1772–1786. doi: 10.1016/j.bbamcr.2012.12.016
- Di Castro, M. A., and Volterra, A. (2022). Astrocyte control of the entorhinal cortex-dentate gyrus circuit: relevance to cognitive processing and impairment in pathology. *Glia* 70, 1536–1553. doi: 10.1002/glia.24128
- Diano, S., Naftolin, F., and Horvath, T. L. (1998). Kainate glutamate receptors (GluR5–7) in the rat arcuate nucleus: relationship to tanycytes, astrocytes, neurons and gonadal steroid receptors. *J. Neuroendocrinol.* 10, 239–247. doi: 10.1046/j.1365-2826.1998.00195.x
- Erecinska, M., Nelson, D., Nissim, I., Daikhin, Y., and Yudkoff, M. (1994). Cerebral alanine transport and alanine aminotransferase reaction: alanine as a source of neuronal glutamate. *J. Neurochem.* 62, 1953–1964. doi: 10.1046/j.1471-4159.1994.62051953.x
- Estrada-Sánchez, A. M., Camacho, A., Montiel, T., and Massieu, L. (2007). Cerebellar granule neurons are more vulnerable to transient transport-mediated glutamate release than to glutamate uptake blockade: correlation with excitatory amino acids levels. *Neurochem. Res.* 32, 423–432. doi: 10.1007/s11064-006-9243-3
- Estrada-Sánchez, A. M., Castro, D., Portillo-Ortiz, K., Jang, K., Nedjat-Haiem, M., Levine, M. S., et al. (2019). Complete but not partial inhibition of glutamate transporters exacerbates cortical excitability in the R6/2 mouse model of Huntington's disease. *CNS Neurosci. Ther.* 25, 509–518. doi: 10.1111/cns.13070
- Estrada-Sánchez, A. M., Levine, M. S., and Cepeda, C. (2017). “Chapter 71 - epilepsy in other neurodegenerative disorders: Huntington's and Parkinson's diseases,” in *Models of Seizures and Epilepsy (Second Edition)*, eds A. Pitkänen, P. S. Buckmaster, A. S. Galanopoulou, and S. L. Moshé (Cambridge, MA: Academic Press), 1043–1058.
- Estrada-Sánchez, A. M., Montiel, T., and Massieu, L. (2010). Glycolysis inhibition decreases the levels of glutamate transporters and enhances glutamate neurotoxicity in the R6/2 Huntington's disease mice. *Neurochem. Res.* 35, 1156–1163. doi: 10.1007/s11064-010-0168-5
- Estrada-Sánchez, A. M., Montiel, T., Segovia, J., and Massieu, L. (2009). Glutamate toxicity in the striatum of the R6/2 Huntington's disease transgenic mice is age-dependent and correlates with decreased levels of glutamate transporters. *Neurobiol. Dis.* 34, 78–86. doi: 10.1016/j.nbd.2008.12.017
- Estrada-Sánchez, A. M., and Rebec, G. V. (2012). Corticostriatal dysfunction and glutamate transporter 1 (GLT1) in Huntington's disease: interactions between neurons and astrocytes. *Basal Ganglia* 2, 57–66. doi: 10.1016/j.baga.2012.04.029
- Faudeau, M., Kim, J., Cormier, K., Gilmore, R., Welch, M., Auregan, G., et al. (2010). *In vivo* expression of polyglutamine-expanded huntingtin by mouse striatal astrocytes impairs glutamate transport: a correlation with Huntington's disease subjects. *Hum. Mol. Genet.* 19, 3053–3067. doi: 10.1093/hmg/dq212
- Fairman, W. A., Vandenberg, R. J., Arriza, J. L., Kavanaugh, M. P., and Amara, S. G. (1995). An excitatory amino-acid transporter with properties of a ligand-gated chloride channel. *Nature* 375, 599–603. doi: 10.1038/375599a0
- Fasano, C., Rocchetti, J., Pietrajitis, K., Zander, J. F., Manseau, F., Sakae, D. Y., et al. (2017). Regulation of the hippocampal network by VGLUT3-positive CCK-GABAergic basket cells. *Front. Cell. Neurosci.* 11:140. doi: 10.3389/fncel.2017.00140
- Fernandes, H. B., Catches, J. S., Petralia, R. S., Copits, B. A., Xu, J., Russell, T. A., et al. (2009). High-affinity kainate receptor subunits are necessary for ionotropic but not metabotropic signaling. *Neuron* 63, 818–829. doi: 10.1016/j.neuron.2009.08.010
- Fernstrom, J. D. (2018). Monosodium glutamate in the diet does not raise brain glutamate concentrations or disrupt brain functions. *Ann. Nutr. Metab.* 73, 43–52. doi: 10.1159/000494782
- Fiacco, T. A., and McCarthy, K. D. (2004). Intracellular astrocyte calcium waves *in situ* increase the frequency of spontaneous AMPA receptor currents in CA1 pyramidal neurons. *J. Neurosci.* 24, 722–732. doi: 10.1523/JNEUROSCI.2859-03.2004
- Freneau, R. T., Jr., Burman, J., Qureshi, T., Tran, C. H., Proctor, J., Johnson, J., et al. (2002). The identification of vesicular glutamate transporter 3 suggests novel modes of signaling by glutamate. *Proc. Natl. Acad. Sci. U S A* 99, 14488–14493. doi: 10.1073/pnas.222546799
- Freneau, R. T., Jr., Troyer, M. D., Pahner, I., Nygaard, G. O., Tran, C. H., Reimer, R. J., et al. (2001). The expression of vesicular glutamate transporters defines two classes of excitatory synapse. *Neuron* 31, 247–260. doi: 10.1016/s0896-6273(01)00344-0
- Fujiyama, F., Furuta, T., and Kaneko, T. (2001). Immunocytochemical localization of candidates for vesicular glutamate transporters in the rat cerebral cortex. *J. Comp. Neurol.* 435, 379–387. doi: 10.1002/cne.1037
- Gegelashvili, G., Dehnes, Y., Danbolt, N. C., and Schousboe, A. (2000). The high-affinity glutamate transporters GLT1, GLAST and EAAT4 are regulated via different signalling mechanisms. *Neurochem. Int.* 37, 163–170. doi: 10.1016/s0197-0186(00)0019-x
- Giaume, C., Tabernero, A., and Medina, J. M. (1997). Metabolic trafficking through astrocytic gap junctions. *Glia* 21, 114–123. doi: 10.1002/(SICI)1098-1136(199709)21:1%3C114::AID-GLIA13%3E3.0.CO;2-V
- Gomez-Gonzalo, M., Martin-Fernandez, M., Martinez-Murillo, R., Mederos, S., Hernandez-Vivanco, A., Jamison, S., et al. (2017). Neuron-astrocyte signaling is preserved in the aging brain. *Glia* 65, 569–580. doi: 10.1002/glia.23112
- Gomez-Gonzalo, M., Zehnder, T., Reque, L. M., Bezzi, P., and Carmignoto, G. (2018). Insights into the release mechanism of astrocytic glutamate evoking in neurons NMDA receptor-mediated slow depolarizing inward currents. *Glia* 66, 2188–2199. doi: 10.1002/glia.23473
- Guerra-Gomes, S., Sousa, N., Pinto, L., and Oliveira, J. F. (2018). Functional roles of astrocyte calcium elevations: from synapses to behavior. *Front. Cell. Neurosci.* 11:427. doi: 10.3389/fncel.2017.00427
- Hamberger, A. C., Chiang, G. H., Nysten, E. S., Scheff, S. W., and Cotman, C. W. (1979). Glutamate as a CNS transmitter. I. Evaluation of glucose and glutamine as precursors for the synthesis of preferentially released glutamate. *Brain Res.* 168, 513–530. doi: 10.1016/0006-8993(79)90306-8
- Hamilton, N. B., and Attwell, D. (2010). Do astrocytes really exocytose neurotransmitters? *Nat. Rev. Neurosci.* 11, 227–238. doi: 10.1038/nrn2803
- Han, J., Kesner, P., Metna-Laurent, M., Duan, T., Xu, L., Georges, F., et al. (2012). Acute cannabinoids impair working memory through astroglial CB1 receptor modulation of hippocampal LTD. *Cell* 148, 1039–1050. doi: 10.1016/j.cell.2012.01.037

- Han, J., Pluhackova, K., and Bockmann, R. A. (2017). The multifaceted role of SNARE proteins in membrane fusion. *Front. Physiol.* 8:5. doi: 10.3389/fphys.2017.00005
- Hansen, K. B., Wollmuth, L. P., Bowie, D., Furukawa, H., Menniti, F. S., Sobolevsky, A. I., et al. (2021). Structure, function and pharmacology of glutamate receptor ion channels. *Pharmacol. Rev.* 73, 298–487. doi: 10.1124/pharmrev.120.000131
- Hardingham, G. E., and Bading, H. (2010). Synaptic versus extrasynaptic NMDA receptor signalling: implications for neurodegenerative disorders. *Nat. Rev. Neurosci.* 11, 682–696. doi: 10.1038/nrn2911
- Hassel, B. (2001). Pyruvate carboxylation in neurons. *J. Neurosci. Res.* 66, 755–762. doi: 10.1002/jnr.10044
- Hawkins, R. A. (2009). The blood-brain barrier and glutamate. *Am. J. Clin. Nutr.* 90, 867S–874S. doi: 10.3945/ajcn.2009.27462BB
- He, L., Vasilou, K., and Nebert, D. W. (2009). Analysis and update of the human solute carrier (SLC) gene superfamily. *Hum. Genomics* 3, 195–206. doi: 10.1186/1479-7364-3-2-195
- Heller, J. P., Odii, T., Zheng, K., and Rusakov, D. A. (2020). Imaging tripartite synapses using super-resolution microscopy. *Methods* 174, 81–90. doi: 10.1016/j.ymeth.2019.05.024
- Henneberger, C., Papouin, T., Oliet, S. H., and Rusakov, D. A. (2010). Long-term potentiation depends on release of D-serine from astrocytes. *Nature* 463, 232–236. doi: 10.1038/nature08673
- Herb, A., Burnashev, N., Werner, P., Sakmann, B., Wisden, W., and Seeburg, P. H. (1992). The KA-2 subunit of excitatory amino acid receptors shows widespread expression in brain and forms ion channels with distantly related subunits. *Neuron* 8, 775–785. doi: 10.1016/0896-6273(92)90098-x
- Herman, M. A., Ackermann, F., Trimbuch, T., and Rosenmund, C. (2014). Vesicular glutamate transporter expression level affects synaptic vesicle release probability at hippocampal synapses in culture. *J. Neurosci.* 34, 11781–11791. doi: 10.1523/JNEUROSCI.1444-14.2014
- Herzog, E., Gilchrist, J., Gras, C., Muzerelle, A., Ravassard, P., Giros, B., et al. (2004). Localization of VGLUT3, the vesicular glutamate transporter type 3, in the rat brain. *Neuroscience* 123, 983–1002. doi: 10.1016/j.neuroscience.2003.10.039
- Hogstad, S., Svenneby, G., Torgner, I. A., Kvamme, E., Hertz, L., and Schousboe, A. (1988). Glutaminase in neurons and astrocytes cultured from mouse brain: kinetic properties and effects of phosphate, glutamate and ammonia. *Neurochem. Res.* 13, 383–388. doi: 10.1007/BF00972489
- Hrabetova, S., Serrano, P., Blace, N., Tse, H. W., Skifter, D. A., Jane, D. E., et al. (2000). Distinct NMDA receptor subpopulations contribute to long-term potentiation and long-term depression induction. *J. Neurosci.* 20:RC81. doi: 10.1523/JNEUROSCI.20-12-j0002.2000
- Huntley, G. W., Rogers, S. W., Moran, T., Janssen, W., Archin, N., Vickers, J. C., et al. (1993). Selective distribution of kainate receptor subunit immunoreactivity in monkey neocortex revealed by a monoclonal antibody that recognizes glutamate receptor subunits GluR5/6/7. *J. Neurosci.* 13, 2965–2981. doi: 10.1523/JNEUROSCI.13-07-02965.1993
- Huyghe, D., Nakamura, Y., Terunuma, M., Faideau, M., Haydon, P., Pangalos, M. N., et al. (2014). Glutamine synthetase stability and subcellular distribution in astrocytes are regulated by γ -aminobutyric type B receptors. *J. Biol. Chem.* 289, 28808–28815. doi: 10.1074/jbc.M114.583534
- Intson, K., Geissah, S., McCullumsmith, R. E., and Ramsey, A. J. (2022). A role for endothelial NMDA receptors in the pathophysiology of schizophrenia. *Schizophr. Res.* 249, 63–73. doi: 10.1016/j.schres.2020.10.004
- Islam, M. M., Nautiyal, M., Wynn, R. M., Mobley, J. A., Chuang, D. T., and Hutson, S. M. (2010). Branched-chain amino acid metabolon: interaction of glutamate dehydrogenase with the mitochondrial branched-chain aminotransferase (BCATm). *J. Biol. Chem.* 285, 265–276. doi: 10.1074/jbc.M109.048777
- Jabaudon, D., Shimamoto, K., Yasuda-Kamatani, Y., Scanziani, M., Gahwiler, B. H., and Gerber, U. (1999). Inhibition of uptake unmasks rapid extracellular turnover of glutamate of nonvesicular origin. *Proc. Natl. Acad. Sci. U S A* 96, 8733–8738. doi: 10.1073/pnas.96.15.8733
- Jeftinija, S. D., Jeftinija, K. V., and Stefanovic, G. (1997). Cultured astrocytes express proteins involved in vesicular glutamate release. *Brain Res.* 750, 41–47. doi: 10.1016/s0006-8993(96)00610-5
- Jimenez-Blasco, D., Santofimia-Castano, P., Gonzalez, A., Almeida, A., and Bolanos, J. P. (2015). Astrocyte NMDA receptors' activity sustains neuronal survival through a Cdk5-Nrf2 pathway. *Cell Death Differ.* 22, 1877–1889. doi: 10.1038/cdd.2015.49
- Juge, N., Yoshida, Y., Yatsushiro, S., Omote, H., and Moriyama, Y. (2006). Vesicular glutamate transporter contains two independent transport machineries. *J. Biol. Chem.* 281, 39499–39506. doi: 10.1074/jbc.M607670200
- Kanai, Y., Bhidé, P. G., DiFiglia, M., and Hediger, M. A. (1995). Neuronal high-affinity glutamate transport in the rat central nervous system. *Neuroreport* 6, 2357–2362. doi: 10.1097/00001756-199511270-00020
- Kanai, Y., and Hediger, M. A. (1992). Primary structure and functional characterization of a high-affinity glutamate transporter. *Nature* 360, 467–471. doi: 10.1038/360467a0
- Kang, J., Jiang, L., Goldman, S. A., and Nedergaard, M. (1998). Astrocyte-mediated potentiation of inhibitory synaptic transmission. *Nat. Neurosci.* 1, 683–692. doi: 10.1038/3684
- Kaufman, A. M., Milnerwood, A. J., Sepers, M. D., Coquinco, A., She, K., Wang, L., et al. (2012). Opposing roles of synaptic and extrasynaptic NMDA receptor signaling in cocultured striatal and cortical neurons. *J. Neurosci.* 32, 3992–4003. doi: 10.1523/JNEUROSCI.4129-11.2012
- Kazama, M., Kato, Y., Kakita, A., Noguchi, N., Urano, Y., Masui, K., et al. (2020). Astrocytes release glutamate via cystine/glutamate antiporter upregulated in response to increased oxidative stress related to sporadic amyotrophic lateral sclerosis. *Neuropathology* 40, 587–598. doi: 10.1111/neup.12716
- Koh, W., Park, M., Chun, Y. E., Lee, J., Shim, H. S., Park, M. G., et al. (2021). Astrocytes render memory flexible. *bioRxiv* [Preprint]. doi: 10.1101/2021.03.25.436945
- Koh, W., Park, M., Chun, Y. E., Lee, J., Shim, H. S., Park, M. G., et al. (2022). Astrocytes render memory flexible by releasing D-serine and regulating NMDA receptor tone in the hippocampus. *Biol. Psychiatry* 91, 740–752. doi: 10.1016/j.biopsych.2021.10.012
- Koutsilieri, E., and Riederer, P. (2007). Excitotoxicity and new antiglutamatergic strategies in Parkinson's disease and Alzheimer's disease. *Parkinsonism Relat. Disord.* 13, S329–S331. doi: 10.1016/S1353-8020(08)70025-7
- Kugler, P., and Schleyer, V. (2004). Developmental expression of glutamate transporters and glutamate dehydrogenase in astrocytes of the postnatal rat hippocampus. *Hippocampus* 14, 975–985. doi: 10.1002/hipo.20015
- Kukley, M., Barden, J. A., Steinhilber, C., and Jabs, R. (2001). Distribution of P2X receptors on astrocytes in juvenile rat hippocampus. *Glia* 36, 11–21. doi: 10.1002/glia.1091
- Kvamme, E., Torgner, I. A., and Roberg, B. (2001). Kinetics and localization of brain phosphate activated glutaminase. *J. Neurosci. Res.* 66, 951–958. doi: 10.1002/jnr.10041
- Kvist, T., Steffensen, T. B., Greenwood, J. R., Mehrzad Tabrizi, F., Hansen, K. B., Gajhede, M., et al. (2013). Crystal structure and pharmacological characterization of a novel N-methyl-D-aspartate (NMDA) receptor antagonist at the GluN1 glycine binding site. *J. Biol. Chem.* 288, 33124–33135. doi: 10.1074/jbc.M113.480210
- Laake, J. H., Takumi, Y., Eidet, J., Torgner, I. A., Roberg, B., Kvamme, E., et al. (1999). Postembedding immunogold labelling reveals subcellular localization and pathway-specific enrichment of phosphate activated glutaminase in rat cerebellum. *Neuroscience* 88, 1137–1151. doi: 10.1016/s0306-4522(98)00298-x
- Lakhan, S. E., Caro, M., and Hadzimechalis, N. (2013). NMDA receptor activity in neuropsychiatric disorders. *Front. Psychiatry* 4:52. doi: 10.3389/fpsy.2013.00052
- Lalo, U., Rasooli-Nejad, S., Bogdanov, A., More, L., Koh, W., Muller, J., et al. (2021). Synergy between vesicular and non-vesicular gliotransmission regulates synaptic plasticity and working memory. *bioRxiv* [Preprint]. doi: 10.1101/2021.03.25.437028
- Lapato, A. S., and Tiwari-Woodruff, S. K. (2018). Connexins and pannexins: at the junction of neuro-glial homeostasis & disease. *J. Neurosci. Res.* 96, 31–44. doi: 10.1002/jnr.24088
- Lauri, S. E., Vesikansa, A., Segerstrale, M., Collingridge, G. L., Isaac, J. T., and Taira, T. (2006). Functional maturation of CA1 synapses involves activity-dependent loss of tonic kainate receptor-mediated inhibition of glutamate release. *Neuron* 50, 415–429. doi: 10.1016/j.neuron.2006.03.020
- Le Meur, K., Galante, M., Angulo, M. C., and Audinat, E. (2007). Tonic activation of NMDA receptors by ambient glutamate of non-synaptic origin in the rat hippocampus. *J. Physiol.* 580, 373–383. doi: 10.1113/jphysiol.2006.123570
- Lee, M., Ko, D. G., Hong, D. K., Lim, M. S., Choi, B. Y., and Suh, S. W. (2020). Role of excitatory amino acid carrier 1 (EAAC1) in neuronal death and neurogenesis after ischemic stroke. *Int. J. Mol. Sci.* 21:5676. doi: 10.3390/ijms21115676
- Lee, M. L., Martinez-Lozada, Z., Krizman, E. N., and Robinson, M. B. (2017). Brain endothelial cells induce astrocytic expression of the glutamate transporter GLT-1 by a Notch-dependent mechanism. *J. Neurochem.* 143, 489–506. doi: 10.1111/jnc.14135
- Lee, M., McGeer, E. G., and McGeer, P. L. (2011). Mechanisms of GABA release from human astrocytes. *Glia* 59, 1600–1611. doi: 10.1002/glia.21202
- Lehre, K. P., Levy, L. M., Ottersen, O. P., Storm-Mathisen, J., and Danbolt, N. C. (1995). Differential expression of two glial glutamate transporters in the rat brain: quantitative and immunocytochemical observations. *J. Neurosci.* 15, 1835–1853. doi: 10.1523/JNEUROSCI.15-03-01835.1995
- Levy, L. M., Lehre, K. P., Rolstad, B., and Danbolt, N. C. (1993). A monoclonal antibody raised against an $[Na^+K^+]$ -coupled L-glutamate transporter purified from rat brain confirms glial cell localization. *FEBS Lett.* 317, 79–84. doi: 10.1016/0014-5793(93)81495-1
- Levy, L. M., Warr, O., and Attwell, D. (1998). Stoichiometry of the glial glutamate transporter GLT-1 expressed inducibly in a Chinese hamster ovary cell line selected for low endogenous Na^+ -dependent glutamate uptake. *J. Neurosci.* 18, 9620–9628. doi: 10.1523/JNEUROSCI.18-23-09620.1998
- Lewerenz, J., Maher, P., and Methner, A. (2012). Regulation of xCT expression and system x(c) function in neuronal cells. *Amino Acids* 42, 171–179. doi: 10.1007/s00726-011-0862-x

- Li, D., Herault, K., Silm, K., Evrard, A., Wojcik, S., Oheim, M., et al. (2013a). Lack of evidence for vesicular glutamate transporter expression in mouse astrocytes. *J. Neurosci.* 33, 4434–4455. doi: 10.1523/JNEUROSCI.3667-12.2013
- Li, D., Agulhon, C., Schmidt, E., Oheim, M., and Ropert, N. (2013b). New tools for investigating astrocyte-to-neuron communication. *Front. Cell. Neurosci.* 7:193. doi: 10.3389/fncel.2013.00193
- Liu, Q. S., Xu, Q., Arcuino, G., Kang, J., and Nedergaard, M. (2004). Astrocyte-mediated activation of neuronal kainate receptors. *Proc. Natl. Acad. Sci. U S A* 101, 3172–3177. doi: 10.1073/pnas.0306731101
- Luque, J. M., and Richards, J. G. (1995). Expression of NMDA 2B receptor subunit mRNA in Bergmann glia. *Glia* 13, 228–232. doi: 10.1002/glia.440130309
- Lyon, K. A., and Allen, N. J. (2022). From synapses to circuits, astrocytes regulate behavior. *Front. Neural Circuits* 15:786293. doi: 10.3389/fncir.2021.786293
- Magi, S., Piccirillo, S., and Amoroso, S. (2019). The dual face of glutamate: from a neurotoxin to a potential survival factor—metabolic implications in health and disease. *Cell. Mol. Life Sci.* 76, 1473–1488. doi: 10.1007/s00018-018-3002-x
- Marcaggi, P., Mutoh, H., Dimitrov, D., Beato, M., and Knopfel, T. (2009). Optical measurement of mGluR1 conformational changes reveals fast activation, slow deactivation and sensitization. *Proc. Natl. Acad. Sci. U S A* 106, 11388–11393. doi: 10.1073/pnas.0901290106
- Martin, R., Bajo-Graneras, R., Moratalla, R., Perea, G., and Araque, A. (2015). Circuit-specific signaling in astrocyte-neuron networks in basal ganglia pathways. *Science* 349, 730–734. doi: 10.1126/science.aaa7945
- Masliah, E., Alford, M., Mallory, M., Rockenstein, E., Moechars, D., and Van Leuven, F. (2000). Abnormal glutamate transport function in mutant amyloid precursor protein transgenic mice. *Exp. Neurol.* 163, 381–387. doi: 10.1006/exnr.2000.7386
- Massie, A., Cnops, L., Smolders, I., McCullumsmith, R., Kooijman, R., Kwak, S., et al. (2008). High-affinity Na⁺/K⁺-dependent glutamate transporter EAAT4 is expressed throughout the rat fore- and midbrain. *J. Comp. Neurol.* 511, 155–172. doi: 10.1002/cne.21823
- Masu, M., Tanabe, Y., Tsuchida, K., Shigemoto, R., and Nakanishi, S. (1991). Sequence and expression of a metabotropic glutamate receptor. *Nature* 349, 760–765. doi: 10.1038/349760a0
- Masuoka, T., Ikeda, R., and Konishi, S. (2019). Persistent activation of histamine H1 receptors in the hippocampal CA1 region enhances NMDA receptor-mediated synaptic excitation and long-term potentiation in astrocyte- and D-serine-dependent manner. *Neuropharmacology* 151, 64–73. doi: 10.1016/j.neuropharm.2019.03.036
- Matschke, V., Theiss, C., Hollmann, M., Schulze-Bahr, E., Lang, F., Seebohm, G., et al. (2015). NDRG2 phosphorylation provides negative feedback for SGK1-dependent regulation of a kainate receptor in astrocytes. *Front. Cell. Neurosci.* 9:387. doi: 10.3389/fncel.2015.00387
- Matthias, K., Kirchhoff, F., Seifert, G., Huttmann, K., Matyash, M., Kettenmann, H., et al. (2003). Segregated expression of AMPA-type glutamate receptors and glutamate transporters defines distinct astrocyte populations in the mouse hippocampus. *J. Neurosci.* 23, 1750–1758. doi: 10.1523/JNEUROSCI.23-05-01750.2003
- McKenna, M. C., Hopkins, I. B., Lindauer, S. L., and Bamford, P. (2006). Aspartate aminotransferase in synaptic and nonsynaptic mitochondria: differential effect of compounds that influence transient hetero-enzyme complex (metabolon) formation. *Neurochem. Int.* 48, 629–636. doi: 10.1016/j.neuint.2005.11.018
- McKenna, M. C., Tildon, J. T., Stevenson, J. H., Huang, X., and Kingwell, K. G. (1995). Regulation of mitochondrial and cytosolic malic enzymes from cultured rat brain astrocytes. *Neurochem. Res.* 20, 1491–1501. doi: 10.1007/BF00970599
- McNair, L. F., Andersen, J. V., Aldana, B. I., Hohnholt, M. C., Nissen, J. D., Sun, Y., et al. (2019). Deletion of neuronal GLT-1 in mice reveals its role in synaptic glutamate homeostasis and mitochondrial function. *J. Neurosci.* 39, 4847–4863. doi: 10.1523/JNEUROSCI.0894-18.2019
- Mennerick, S., Dhond, R. P., Benz, A., Xu, W., Rothstein, J. D., Danbolt, N. C., et al. (1998). Neuronal expression of the glutamate transporter GLT-1 in hippocampal microcultures. *J. Neurosci.* 18, 4490–4499. doi: 10.1523/JNEUROSCI.18-12-04490.1998
- Meyerson, J. R., Chittori, S., Merk, A., Rao, P., Han, T. H., Serpe, M., et al. (2016). Structural basis of kainate subtype glutamate receptor desensitization. *Nature* 537, 567–571. doi: 10.1038/nature19352
- Milnerwood, A. J., Kaufman, A. M., Sepers, M. D., Gladding, C. M., Zhang, L., Wang, L., et al. (2012). Mitigation of augmented extrasynaptic NMDAR signaling and apoptosis in cortico-striatal co-cultures from Huntington's disease mice. *Neurobiol. Dis.* 48, 40–51. doi: 10.1016/j.nbd.2012.05.013
- Min, R., and Nevian, T. (2012). Astrocyte signaling controls spike timing-dependent depression at neocortical synapses. *Nat. Neurosci.* 15, 746–753. doi: 10.1038/nn.3075
- Mineff, E., and Valtschanoff, J. (1999). Metabotropic glutamate receptors 2 and 3 expressed by astrocytes in rat ventrobasal thalamus. *Neurosci. Lett.* 270, 95–98. doi: 10.1016/s0304-3940(99)00484-x
- Montana, V., Ni, Y., Sunjara, V., Hua, X., and Parpura, V. (2004). Vesicular glutamate transporter-dependent glutamate release from astrocytes. *J. Neurosci.* 24, 2633–2642. doi: 10.1523/JNEUROSCI.3770-03.2004
- Montero, T. D., and Orellana, J. A. (2015). Hemichannels: new pathways for gliotransmitter release. *Neuroscience* 286, 45–59. doi: 10.1016/j.neuroscience.2014.11.048
- Mothet, J. P., Pollegioni, L., Ouanounou, G., Martineau, M., Fossier, P., and Baux, G. (2005). Glutamate receptor activation triggers a calcium-dependent and SNARE protein-dependent release of the gliotransmitter D-serine. *Proc. Natl. Acad. Sci. U S A* 102, 5606–5611. doi: 10.1073/pnas.0408483102
- Müller, T., Möller, T., Berger, T., Schnitzer, J., and Kettenmann, H. (1992). Calcium entry through kainate receptors and resulting potassium-channel blockade in Bergmann glial cells. *Science* 256, 1563–1566. doi: 10.1126/science.1317969
- Nakamura, E., Sato, M., Yang, H., Miyagawa, F., Harasaki, M., Tomita, K., et al. (1999). 4F2 (CD98) heavy chain is associated covalently with an amino acid transporter and controls intracellular trafficking and membrane topology of 4F2 heterodimer. *J. Biol. Chem.* 274, 3009–3016. doi: 10.1074/jbc.274.5.3009
- Navarrete, M., Perea, G., Maglio, L., Pastor, J., Garcia de Sola, R., and Araque, A. (2013). Astrocyte calcium signal and gliotransmission in human brain tissue. *Cereb. Cortex* 23, 1240–1246. doi: 10.1093/cercor/bhs122
- Neame, S., Safory, H., Radziszewsky, I., Toutou, A., Marchesani, F., Marchetti, M., et al. (2019). The NMDA receptor activation by d-serine and glycine is controlled by an astrocytic Phgdh-dependent serine shuttle. *Proc. Natl. Acad. Sci. U S A* 116, 20736–20742. doi: 10.1073/pnas.1909458116
- Nedergaard, M. (1994). Direct signaling from astrocytes to neurons in cultures of mammalian brain cells. *Science* 263, 1768–1771. doi: 10.1126/science.8134839
- Newman, E. A., and Zahs, K. R. (1998). Modulation of neuronal activity by glial cells in the retina. *J. Neurosci.* 18, 4022–4028. doi: 10.1523/JNEUROSCI.18-11-04022.1998
- Norenberg, M. D., and Martinez-Hernandez, A. (1979). Fine structural localization of glutamine synthetase in astrocytes of rat brain. *Brain Res.* 161, 303–310. doi: 10.1016/0006-8993(79)90071-4
- O'Donovan, S. M., Sullivan, C. R., and McCullumsmith, R. E. (2017). The role of glutamate transporters in the pathophysiology of neuropsychiatric disorders. *NPJ Schizophr.* 3:32. doi: 10.1038/s41537-017-0037-1
- Ong, W. Y., Tanaka, K., Dawe, G. S., Ittner, L. M., and Farooqui, A. A. (2013). Slow excitotoxicity in Alzheimer's disease. *J. Alzheimers Dis.* 35, 643–668. doi: 10.3233/JAD-121990
- Ormel, L., Stensrud, M. J., Bergersen, L. H., and Gundersen, V. (2012). VGLUT1 is localized in astrocytic processes in several brain regions. *Glia* 60, 229–238. doi: 10.1002/glia.21258
- Osterberg, K. A., and Wattenberg, L. W. (1962). Oxidative histochemistry of reactive astrocytes. *Arch. Neurol.* 7, 211–218. doi: 10.1001/archneur.1962.04210030049007
- Ottstad-Hansen, S., Hu, Q. X., Follin-Arbelet, V. V., Bentea, E., Sato, H., Massie, A., et al. (2018). The cystine-glutamate exchanger (xCT, Slc7a11) is expressed in significant concentrations in a subpopulation of astrocytes in the mouse brain. *Glia* 66, 951–970. doi: 10.1002/glia.23294
- Pajarillo, E., Rizor, A., Lee, J., Aschner, M., and Lee, E. (2019). The role of astrocytic glutamate transporters GLT-1 and GLAST in neurological disorders: potential targets for neurotherapeutics. *Neuropharmacology* 161:107559. doi: 10.1016/j.neuropharm.2019.03.002
- Palygin, O., Lalo, U., and Pankratov, Y. (2011). Distinct pharmacological and functional properties of NMDA receptors in mouse cortical astrocytes. *Br. J. Pharmacol.* 163, 1755–1766. doi: 10.1111/j.1476-5381.2011.01374.x
- Palygin, O., Lalo, U., Verkhratsky, A., and Pankratov, Y. (2010). Ionotropic NMDA and P2X1/5 receptors mediate synaptically induced Ca²⁺ signalling in cortical astrocytes. *Cell Calcium* 48, 225–231. doi: 10.1016/j.cecc.2010.09.004
- Perea, G., and Araque, A. (2007). Astrocytes potentiate transmitter release at single hippocampal synapses. *Science* 317, 1083–1086. doi: 10.1126/science.1144640
- Perea, G., Gomez, R., Mederos, S., Covelo, A., Ballesteros, J. J., Schlosser, L., et al. (2016). Activity-dependent switch of GABAergic inhibition into glutamatergic excitation in astrocyte-neuron networks. *eLife* 5:e20362. doi: 10.7554/eLife.20362
- Perea, G., Yang, A., Boyden, E. S., and Sur, M. (2014). Optogenetic astrocyte activation modulates response selectivity of visual cortex neurons *in vivo*. *Nat. Commun.* 5:3262. doi: 10.1038/ncomms4262
- Perego, C., Vanoni, C., Bossi, M., Massari, S., Basudev, H., Longhi, R., et al. (2000). The GLT-1 and GLAST glutamate transporters are expressed on morphologically distinct astrocytes and regulated by neuronal activity in primary hippocampal cocultures. *J. Neurochem.* 75, 1076–1084. doi: 10.1046/j.1471-4159.2000.0751076.x
- Peterson, A. R., and Binder, D. K. (2019). Regulation of synaptosomal GLT-1 and GLAST during epileptogenesis. *Neuroscience* 411, 185–201. doi: 10.1016/j.neuroscience.2019.05.048
- Petr, G. T., Sun, Y., Frederick, N. M., Zhou, Y., Dhamne, S. C., Hameed, M. Q., et al. (2015). Conditional deletion of the glutamate transporter GLT-1 reveals that astrocytic GLT-1 protects against fatal epilepsy while neuronal GLT-1 contributes significantly to glutamate uptake into synaptosomes. *J. Neurosci.* 35, 5187–5201. doi: 10.1523/JNEUROSCI.4255-14.2015
- Pines, G., Danbolt, N. C., Bjoras, M., Zhang, Y., Bendahan, A., Eide, L., et al. (1992). Cloning and expression of a rat brain L-glutamate transporter. *Nature* 360, 464–467. doi: 10.1038/360464a0

- Pirttimäki, T. M., Hall, S. D., and Parri, H. R. (2011). Sustained neuronal activity generated by glial plasticity. *J. Neurosci.* 31, 7637–7647. doi: 10.1523/JNEUROSCI.5783-10.2011
- Plaitakis, A., Kalef-Ezra, E., Kotzamani, D., Zaganas, I., and Spanaki, C. (2017). The glutamate dehydrogenase pathway and its roles in cell and tissue biology in health and disease. *Biology (Basel)* 6:11. doi: 10.3390/biology6010011
- Radulescu, A. R., Todd, G. C., Williams, C. L., Bennink, B. A., Lemus, A. A., Chesbro, H. E., et al. (2022). Estimating the glutamate transporter surface density in distinct sub-cellular compartments of mouse hippocampal astrocytes. *PLoS Comput. Biol.* 18:e1009845. doi: 10.1371/journal.pcbi.1009845
- Ravikrishnan, A., Gandhi, P. J., Shelkar, G. P., Liu, J., Pavuluri, R., and Dravid, S. M. (2018). Region-specific expression of NMDA receptor GluN2C subunit in parvalbumin-positive neurons and astrocytes: analysis of GluN2C expression using a novel reporter model. *Neuroscience* 380, 49–62. doi: 10.1016/j.neuroscience.2018.03.011
- Re, D. B., Nafia, I., Melon, C., Shimamoto, K., Kerkerian-Le Goff, L., and Had-Aissouni, L. (2006). Glutamate leakage from a compartmentalized intracellular metabolic pool and activation of the lipoxygenase pathway mediate oxidative astrocyte death by reversed glutamate transport. *Glia* 54, 47–57. doi: 10.1002/glia.20353
- Reimers, J. M., Milovanovic, M., and Wolf, M. E. (2011). Quantitative analysis of AMPA receptor subunit composition in addiction-related brain regions. *Brain Res.* 1367, 223–233. doi: 10.1016/j.brainres.2010.10.016
- Reiner, A., and Levitz, J. (2018). Glutamatergic signaling in the central nervous system: ionotropic and metabotropic receptors in concert. *Neuron* 98, 1080–1098. doi: 10.1016/j.neuron.2018.05.018
- Rodriguez-Prados, M., Rojo-Ruiz, J., Garcia-Sancho, J., and Alonso, M. T. (2020). Direct monitoring of ER Ca²⁺ dynamics reveals that Ca²⁺ entry induces ER-Ca²⁺ release in astrocytes. *Pflugers Arch.* 472, 439–448. doi: 10.1007/s00424-020-02364-7
- Rothstein, J. D., Dykes-Hoberg, M., Pardo, C. A., Bristol, L. A., Jin, L., Kuncl, R. W., et al. (1996). Knockout of glutamate transporters reveals a major role for astroglial transport in excitotoxicity and clearance of glutamate. *Neuron* 16, 675–686. doi: 10.1016/s0896-6273(00)80086-0
- Rudolph, R., Jahn, H. M., Courjaret, R., Messemer, N., Kirchhoff, F., and Deitmer, J. W. (2016). The inhibitory input to the medial entorhinal Purkinje cells is reciprocally modulated by Bergmann glial P2Y1 and AMPA receptor signaling. *Glia* 64, 1265–1280. doi: 10.1002/glia.22999
- Ruscak, M., Orlicky, J., Zubor, V., and Hager, H. (1982). Alanine aminotransferase in bovine brain: purification and properties. *J. Neurochem.* 39, 210–216. doi: 10.1111/j.1471-4159.1982.tb04720.x
- Sahlender, D. A., Savtchouk, I., and Volterra, A. (2014). What do we know about gliotransmitter release from astrocytes? *Philos. Trans. R Soc. Lond B Biol. Sci.* 369:20130592. doi: 10.1098/rstb.2013.0592
- Salas, I. H., Burgado, J., and Allen, N. J. (2020). Glia: victims or villains of the aging brain? *Neurobiol. Dis.* 143:105008. doi: 10.1016/j.nbd.2020.105008
- Santello, M., Bezzi, P., and Volterra, A. (2011). TNF α controls glutamatergic gliotransmission in the hippocampal dentate gyrus. *Neuron* 69, 988–1001. doi: 10.1016/j.neuron.2011.02.003
- Sato, H., Tamba, M., Ishii, T., and Bannai, S. (1999). Cloning and expression of a plasma membrane cystine/glutamate exchange transporter composed of two distinct proteins. *J. Biol. Chem.* 274, 11455–11458. doi: 10.1074/jbc.274.17.11455
- Sato, H., Tamba, M., Okuno, S., Sato, K., Keino-Masu, K., Masu, M., et al. (2002). Distribution of cystine/glutamate exchange transporter, system x(c)⁻, in the mouse brain. *J. Neurosci.* 22, 8028–8033. doi: 10.1523/JNEUROSCI.22-18-08028.2002
- Savtchouk, I., Di Castro, M. A., Ali, R., Stubbe, H., Lujan, R., and Volterra, A. (2019). Circuit-specific control of the medial entorhinal inputs to the dentate gyrus by atypical presynaptic NMDARs activated by astrocytes. *Proc. Natl. Acad. Sci. U S A* 116, 13602–13610. doi: 10.1073/pnas.1816013116
- Savtchouk, I., and Volterra, A. (2018). Gliotransmission: beyond black-and-white. *J. Neurosci.* 38, 14–25. doi: 10.1523/JNEUROSCI.0017-17.2017
- Schools, G. P., and Kimelberg, H. K. (1999). mGluR3 and mGluR5 are the predominant metabotropic glutamate receptor mRNAs expressed in hippocampal astrocytes acutely isolated from young rats. *J. Neurosci. Res.* 58, 533–543. doi: 10.1002/(sici)1097-4547(19991115)58:4<533::aid-jnr6>3.0.co;2-g
- Schorge, S., Elenes, S., and Colquhoun, D. (2005). Maximum likelihood fitting of single channel NMDA activity with a mechanism composed of independent dimers of subunits. *J. Physiol.* 569, 395–418. doi: 10.1113/jphysiol.2005.095349
- Schousboe, A. (2017). A tribute to Mary C. McKenna: glutamate as energy substrate and neurotransmitter-functional interaction between neurons and astrocytes. *Neurochem. Res.* 42, 4–9. doi: 10.1007/s11064-015-1813-9
- Schousboe, A., Bak, L. K., and Waagepetersen, H. S. (2013). Astrocytic control of biosynthesis and turnover of the neurotransmitters glutamate and GABA. *Front. Endocrinol. (Lausanne)* 4:102. doi: 10.3389/fendo.2013.00102
- Schousboe, A., Scafidi, S., Bak, L. K., Waagepetersen, H. S., and McKenna, M. C. (2014). Glutamate metabolism in the brain focusing on astrocytes. *Adv. Neurobiol.* 11, 13–30. doi: 10.1007/978-3-319-08894-5_2
- Schousboe, A., Waagepetersen, H. S., and Sonnewald, U. (2019). Astrocytic pyruvate carboxylation: status after 35 years. *J. Neurosci. Res.* 97, 890–896. doi: 10.1002/jnr.24402
- Schreiner, A. E., Durry, S., Aida, T., Stock, M. C., Ruther, U., Tanaka, K., et al. (2014). Laminar and subcellular heterogeneity of GLAST and GLT-1 immunoreactivity in the developing postnatal mouse hippocampus. *J. Comp. Neurol.* 522, 204–224. doi: 10.1002/cne.23450
- Schröder, W., Seifert, G., Huttmann, K., Hinterkeuser, S., and Steinhauser, C. (2002). AMPA receptor-mediated modulation of inward rectifier K⁺ channels in astrocytes of mouse hippocampus. *Mol. Cell. Neurosci.* 19, 447–458. doi: 10.1006/mcne.2001.1080
- Schubert, D., and Piasecki, D. (2001). Oxidative glutamate toxicity can be a component of the excitotoxicity cascade. *J. Neurosci.* 21, 7455–7462. doi: 10.1523/JNEUROSCI.21-19-07455.2001
- Seifert, G., Rehn, L., Weber, M., and Steinhäuser, C. (1997). AMPA receptor subunits expressed by single astrocytes in the juvenile mouse hippocampus. *Brain Res. Mol. Brain Res.* 47, 286–294. doi: 10.1016/s0169-328x(97)00059-4
- Seifert, G., and Steinhauser, C. (1995). Glial cells in the mouse hippocampus express AMPA receptors with an intermediate Ca²⁺ permeability. *Eur. J. Neurosci.* 7, 1872–1881. doi: 10.1111/j.1460-9568.1995.tb00708.x
- Shah, P., Plitman, E., Iwata, Y., Kim, J., Nakajima, S., Chan, N., et al. (2020). Glutamatergic neurometabolites and cortical thickness in treatment-resistant schizophrenia: implications for glutamate-mediated excitotoxicity. *J. Psychiatr. Res.* 124, 151–158. doi: 10.1016/j.jpsychires.2020.02.032
- Shashidharan, P., Huntley, G. W., Murray, J. M., Buku, A., Moran, T., Walsh, M. J., et al. (1997). Immunohistochemical localization of the neuron-specific glutamate transporter EAAC1 (EAAT3) in rat brain and spinal cord revealed by a novel monoclonal antibody. *Brain Res.* 773, 139–148. doi: 10.1016/s0006-8993(97)00921-9
- Shen, W., Nikolic, L., Meunier, C., Pfrieger, F., and Audinat, E. (2017). An autocrine purinergic signaling controls astrocyte-induced neuronal excitation. *Sci. Rep.* 7:11280. doi: 10.1038/s41598-017-11793-x
- Sherwood, M. W., Arizono, M., Hisatsune, C., Bannai, H., Ebisui, E., Sherwood, J. L., et al. (2017). Astrocytic IP₃ R: Contribution to Ca²⁺ signalling and hippocampal LTP. *Glia* 65, 502–513. doi: 10.1002/glia.23107
- Sherwood, M. W., Oliet, S. H. R., and Panatier, A. (2021a). NMDARs, coincidence detectors of astrocytic and neuronal activities. *Int. J. Mol. Sci.* 22:7258. doi: 10.3390/ijms22147258
- Sherwood, M. W., Arizono, M., Panatier, A., Mikoshiba, K., and Oliet, S. H. R. (2021b). Astrocytic IP₃Rs: beyond IP₃R2. *Front. Cell. Neurosci.* 15:695817. doi: 10.3389/fncel.2021.695817
- Shibasaki, K., Hosoi, N., Kaneko, R., Tominaga, M., and Yamada, K. (2017). Glycine release from astrocytes via functional reversal of GlyT1. *J. Neurochem.* 140, 395–403. doi: 10.1111/jnc.13741
- Shigetomi, E., Bowser, D. N., Sofroniew, M. V., and Khakh, B. S. (2008). Two forms of astrocyte calcium excitability have distinct effects on NMDA receptor-mediated slow inward currents in pyramidal neurons. *J. Neurosci.* 28, 6659–6663. doi: 10.1523/JNEUROSCI.1717-08.2008
- Shin, C.-S., Mishra, P., Watrous, J. D., Carelli, V., D'Aurelio, M., Jain, M., et al. (2017). The glutamate/cystine xCT antiporter antagonizes glutamine metabolism and reduces nutrient flexibility. *Nat. Commun.* 8:15074. doi: 10.1038/ncomms15074
- Shiratori-Hayashi, M., Yamaguchi, C., Eguchi, K., Shiraiishi, Y., Kohno, K., Mikoshiba, K., et al. (2021). Astrocytic STAT3 activation and chronic itch require IP₃R1/TRPC-dependent Ca²⁺ signals in mice. *J. Allergy Clin. Immunol.* 147, 1341–1353. doi: 10.1016/j.jaci.2020.06.039
- Simard, M., Arcuino, G., Takano, T., Liu, Q. S., and Nedergaard, M. (2003). Signaling at the gliovascular interface. *J. Neurosci.* 23, 9254–9262. doi: 10.1523/JNEUROSCI.23-27-09254.2003
- Sladeczek, F., Pin, J. P., Recasens, M., Bockaert, J., and Weiss, S. (1985). Glutamate stimulates inositol phosphate formation in striatal neurones. *Nature* 317, 717–719. doi: 10.1038/317717a0
- Spampinato, S. F., Copani, A., Nicoletti, F., Sortino, M. A., and Caraci, F. (2018). Metabotropic glutamate receptors in glial cells: a new potential target for neuroprotection? *Front. Mol. Neurosci.* 11:414. doi: 10.3389/fnmol.2018.00414
- Storck, T., Schulte, S., Hofmann, K., and Stoffel, W. (1992). Structure, expression and functional analysis of a Na⁺-dependent glutamate/aspartate transporter from rat brain. *Proc. Natl. Acad. Sci. U S A* 89, 10955–10959. doi: 10.1073/pnas.89.22.10955
- Sugiyama, H., Ito, I., and Hirono, C. (1987). A new type of glutamate receptor linked to inositol phospholipid metabolism. *Nature* 325, 531–533. doi: 10.1038/325531a0
- Sun, W., McConnell, E., Pare, J. F., Xu, Q., Chen, M., Peng, W., et al. (2013). Glutamate-dependent neuroglial calcium signaling differs between young and adult brain. *Science* 339, 197–200. doi: 10.1126/science.1226740
- Swanson, R. A., Liu, J., Miller, J. W., Rothstein, J. D., Farrell, K., Stein, B. A., et al. (1997). Neuronal regulation of glutamate transporter subtype expression in astrocytes. *J. Neurosci.* 17, 932–940. doi: 10.1523/JNEUROSCI.17-03-00932.1997

- Sweeney, P., Qi, Y., Xu, Z., and Yang, Y. (2016). Activation of hypothalamic astrocytes suppresses feeding without altering emotional states. *Glia* 64, 2263–2273. doi: 10.1002/glia.23073
- Takata-Tsuji, F., Chounlamountri, N., Do, L. D., Philippot, C., Novion Ducassou, J., Coute, Y., et al. (2021). Microglia modulate gliotransmission through the regulation of VAMP2 proteins in astrocytes. *Glia* 69, 61–72. doi: 10.1002/glia.23884
- Tanaka, K., Watase, K., Manabe, T., Yamada, K., Watanabe, M., Takahashi, K., et al. (1997). Epilepsy and exacerbation of brain injury in mice lacking the glutamate transporter GLT-1. *Science* 276, 1699–1702. doi: 10.1126/science.276.5319.1699
- Traynelis, S. F., Wollmuth, L. P., McBain, C. J., Menniti, F. S., Vance, K. M., Ogden, K. K., et al. (2010). Glutamate receptor ion channels: structure, regulation and function. *Pharmacol. Rev.* 62, 405–496. doi: 10.1124/pr.109.002451
- Ulbrich, M. H., and Isacoff, E. Y. (2008). Rules of engagement for NMDA receptor subunits. *Proc. Natl. Acad. Sci. U S A* 105, 14163–14168. doi: 10.1073/pnas.0802075105
- Varela, L., Stutz, B., Song, J. E., Kim, J. G., Liu, Z. W., Gao, X. B., et al. (2021). Hunger-promoting AgRP neurons trigger an astrocyte-mediated feed-forward autoactivation loop in mice. *J. Clin. Invest.* 131:e144239. doi: 10.1172/JCI144239
- Vargas, J. R., Takahashi, D. K., Thomson, K. E., and Wilcox, K. S. (2013). The expression of kainate receptor subunits in hippocampal astrocytes after experimentally induced status epilepticus. *J. Neuropathol. Exp. Neurol.* 72, 919–932. doi: 10.1097/NEN.0b013e3182a4b266
- Ventura, R., and Harris, K. M. (1999). Three-dimensional relationships between hippocampal synapses and astrocytes. *J. Neurosci.* 19, 6897–6906. doi: 10.1523/JNEUROSCI.19-16-06897.1999
- Verkhratsky, A., and Kirchhoff, F. (2007). NMDA receptors in glia. *Neuroscientist* 13, 28–37. doi: 10.1177/1073858406294270
- Vermeiren, C., Najimi, M., Vanhoutte, N., Tilleux, S., de Hemptinne, I., Maloteaux, J. M., et al. (2005). Acute up-regulation of glutamate uptake mediated by mGluR5a in reactive astrocytes. *J. Neurochem.* 94, 405–416. doi: 10.1111/j.1471-4159.2005.03216.x
- Waagepetersen, H. S., Sonnewald, U., Larsson, O. M., and Schousboe, A. (2000). A possible role of alanine for ammonia transfer between astrocytes and glutamatergic neurons. *J. Neurochem.* 75, 471–479. doi: 10.1046/j.1471-4159.2000.0750471.x
- Walker, V. (2014). Ammonia metabolism and hyperammonemic disorders. *Adv. Clin. Chem.* 67, 73–150. doi: 10.1016/bs.acc.2014.09.002
- Wang, Q., Kong, Y., Wu, D. Y., Liu, J. H., Jie, W., You, Q. L., et al. (2021). Impaired calcium signaling in astrocytes modulates autism spectrum disorder-like behaviors in mice. *Nat. Commun.* 12:3321. doi: 10.1038/s41467-021-23843-0
- Westergaard, N., Varming, T., Peng, L., Sonnewald, U., Hertz, L., and Schousboe, A. (1993). Uptake, release and metabolism of alanine in neurons and astrocytes in primary cultures. *J. Neurosci. Res.* 35, 540–545. doi: 10.1002/jnr.490350510
- Wiktorska, L., Bilecki, W., Tertilt, M., Kudla, L., Szumiec, L., Mackowiak, M., et al. (2021). Knockdown of the astrocytic glucocorticoid receptor in the central nucleus of the amygdala diminishes conditioned fear expression and anxiety. *Behav. Brain Res.* 402:113095. doi: 10.1016/j.bbr.2020.113095
- Wojcik, S. M., Rhee, J. S., Herzog, E., Sigler, A., Jahn, R., Takamori, S., et al. (2004). An essential role for vesicular glutamate transporter 1 (VGLUT1) in postnatal development and control of quantal size. *Proc. Natl. Acad. Sci. U S A* 101, 7158–7163. doi: 10.1073/pnas.0401764101
- Wolosker, H., de Souza, D. O., and de Meis, L. (1996). Regulation of glutamate transport into synaptic vesicles by chloride and proton gradient. *J. Biol. Chem.* 271, 11726–11731. doi: 10.1074/jbc.271.20.11726
- Woo, D. H., Han, K. S., Shim, J. W., Yoon, B. E., Kim, E., Bae, J. Y., et al. (2012). TREK-1 and Best1 channels mediate fast and slow glutamate release in astrocytes upon GPCR activation. *Cell* 151, 25–40. doi: 10.1016/j.cell.2012.09.005
- Yamada, D., Kawabe, K., Tosa, I., Tsukamoto, S., Nakazato, R., Kou, M., et al. (2019). Inhibition of the glutamine transporter SNAT1 confers neuroprotection in mice by modulating the mTOR-autophagy system. *Commun. Biol.* 2:346. doi: 10.1038/s42003-019-0582-4
- Yu, A. C., Hertz, E., and Hertz, L. (1984). Alterations in uptake and release rates for GABA, glutamate and glutamine during biochemical maturation of highly purified cultures of cerebral cortical neurons, a GABAergic preparation. *J. Neurochem.* 42, 951–960. doi: 10.1111/j.1471-4159.1984.tb12696.x
- Zhang, Q., Fukuda, M., Van Bockstaele, E., Pascual, O., and Haydon, P. G. (2004). Synaptotagmin IV regulates glial glutamate release. *Proc. Natl. Acad. Sci. U S A* 101, 9441–9446. doi: 10.1073/pnas.0401960101
- Zhou, X., Hollern, D., Liao, J., Andrechek, E., and Wang, H. (2013). NMDA receptor-mediated excitotoxicity depends on the coactivation of synaptic and extrasynaptic receptors. *Cell Death Dis.* 4:e560. doi: 10.1038/cddis.2013.82